Stellar Multiplicity

Gaspard Duchêne^{1,2} & Adam Kraus³

¹University of California Berkeley, Hearst Field Annex B-20, Berkeley CA 94720-3411 USA ²UJF-Grenoble 1/ CNRS-INSU, Institut de Planétologie et d'Astrophysique de Grenoble (IPAG) UMR 5274, BP 53, 38041 Grenoble Cedex 9, France ³Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

Key Words

main sequence stars, pre main sequence stars, protostars, star formation, brown dwarfs

Abstract

Stellar multiplicity is an ubiquitous outcome of the star formation process. Characterizing the frequency and main characteristics of multiple systems and their dependencies on primary mass and environment is therefore a powerful tool to probe this process. While early attempts were fraught with selection biases and limited completeness, instrumentation breakthroughs in the last two decades now enable robust analyses. In this review, we summarize our current empirical knowledge of stellar multiplicity for Main Sequence stars and brown dwarfs, as well as among populations of Pre-Main Sequence stars and embedded protostars. Clear trends as a function of both primary mass and stellar evolutionary stage are identified that will serve as a comparison basis for numerical and analytical models of star formation.

1 INTRODUCTION

The existence of binary and multiple systems was recognized soon after the birth of modern astronomy in the 17th century. Kuiper (1935a) first argued that empirically determining the multiplicity frequency and distribution of key orbital parameters would prove to be highly valuable from a theoretical standpoint. The dependency of the multiplicity frequency and their associated orbital parameters on primary mass should contain the imprint of the physical processes at play throughout the lifetime of stellar populations. In particular, if dynamical interactions (within multiple systems, in stellar clusters and in the Galactic tidal field) can be properly accounted for, statistical properties of multiple systems can serve as a posteriori tests for star formation theories. For instance, whether star formation is characterized by universal multiplicity distributions (akin to the apparently-universal Initial Mass Function, IMF, in the present day Universe) is an open question that has important bearing on star formation theory.

The frequency and properties of multiple systems have been discussed in multiple entries in the Annual Review: van de Kamp (1975) discussed astrometric binaries, while Abt (1983) and Mathieu (1994) presented statistical analyses of multiplicity among Main Sequence (MS) field stars in and Pre-Main Sequence (PMS) stars, respectively. Knowledge of stellar multiplicity has greatly expanded since these seminal reviews, due to improvements in instrumentation and larger, more uniform samples. In addition, new stellar populations have been probed, such as high-mass stars (Zinnecker & Yorke 2007) and very low-mass (VLM) stars and brown dwarfs (BDs, see Luhman 2012), allowing for a more complete picture to be drawn. On the theoretical side, prompt fragmentation during the gravitational collapse of a prestellar core appears to be the favored formation mechanism although no single process appears to account for all observed multiplicity properties (Tohline 2002).

This review summarizes our current empirical knowledge of the frequency and principal properties (such as the distributions of orbital periods, mass ratios, and eccentricities) of multiple systems as a function of stellar mass and age. Multiple systems are thought to play central roles in numerous aspects of post-MS evolution, such as the blue straggler phenomenon, X-ray binaries, Type Ia supernovae, and the shaping of bipolar planetary nebulae. However, we restrict the scope of this review to MS and less evolved objects, where unbiased uniform surveys have targeted larger samples.

2 CHARACTERIZING MULTIPLICITY AND SURVEY METHOD-OLOGY

2.1 Key multiplicity properties

The frequency of multiple systems and the companion frequency (average number of companion per target, which can exceed 100%), MF and CF respectively, are the most immediately quantifiable aspect of multiplicity. The difference between these two quantities hinges on the frequency of high-order multiple systems. Unfortunately, these two quantities are frequently confused in the literature, leading to ambiguous interpretations. Beyond these global quantities, and although the diversity of multiple systems is extreme, their statistical properties can be summarized in a handful of distributions, which we briefly introduce here. While correlations between these distributions could be present, multidimensional analyses require very large samples of binaries, which remain out of reach in most cases.

The orbital period, P, of a binary system can be readily estimated for spectroscopic binaries (SBs). For visual binaries (VBs), the orbital semi-major axis, a, can be estimated from the projected separation, ρ , on a statistical basis (e.g., Brandeker et al. 2006). A power law parametrization, $f(P) \propto P^{\alpha}$, is commonly used, with the particular case $\alpha = -1$ (a logarithmically-flat distribution known as "Öpik's law", Öpik 1924) being frequently favored. An alternative log-normal representation, parametrized by \overline{P} and $\sigma_{\log P}$, is also widely used, although it differs only marginally from Öpik's law for large values of $\sigma_{\log P}$ (Heacox 1996). Distinguishing empirically between these two parametrizations would have fundamental implications on the binary formation process, as Öpik's law suggests a scalefree process while a log-normal function implies a preferred spatial scale for companion formation. However, dynamical evolution throughout a system's lifetime can significantly alter its period, complicating the interpretation.

The mass ratio, $q = M_{\text{sec}}/M_{\text{prim}} \leq 1$, can generally be derived from the observed flux ratio, although only a model-dependent estimate can be obtained for PMS and substellar objects. A power law distribution, $f(q) \propto q^{\gamma}$, is frequently used to characterize the distribution of mass ratios and we use this formalism in this review to compare different populations. When no robust estimate of γ has been proposed in the literature, we computed it from available data. The orbital eccentricity is an important tool to probe the dynamical evolution of a system: short period binaries can circularize as a result of tidal dissipation (Koch & Hrivnak 1981), while eccentricity can be pumped in three-body internal (Kozai) and external interactions. The expected distribution for a dynamically relaxed ("thermal") population is f(e) = 2e (Ambartsumian 1937). Because eccentricity can only be reliably estimated for binaries whose orbit is short enough to be fully mapped, only a subset of all systems can be included in the analysis of the eccentricity distribution. We limit our discuss of this quantity to Section 5.1.4. Finally, the other orbital parameters define the orientation of the orbital plane relative to our viewing geometry and are therefore not as astrophysically compelling. The orientation of orbital planes is essentially random with respect to our line of sight and to the stellar rotation spin vector (e.g., Hale 1994; Popovic, Pavlovic & Ninkovic 2004), while multiple systems only have a modest preference for internal coplanarity (e.g., Sterzik & Tokovinin 2002). We will not discuss these quantities further in this review.

2.2 General methodology

An optimal multiplicity survey should 1) rely on a complete, uniform search of a large, volume-limited sample and 2) make use of a variety of observing methods, as each of them is tailored to a certain type of companions. Magnitude-limited surveys, which are very common, are affected by an inevitable Malmquist-like bias, but this effect can be corrected statistically. Still, such surveys can be affected by more subtle selections biases, such as preferences for less extincted and/or more massive (brighter) targets. A sample size of at least 100 targets is typically required to measure multiplicity frequencies with a precision of $\pm 5\%$. Studies of orbital parameters (e.g., P or q) require a similar number of companions, hence a sample that is several times larger.

Whenever possible, the comparison between two independent surveys should be performed directly over a common section of the parameter space, and binomial statistics should be preferred (see Brandeker et al. 2006). However, most multiplicity surveys are incomplete and it is normal practice to estimate a completeness correction to take "missed companions" into account. Bayesian inference techniques can also be used to optimally exploit incomplete data while properly presenting the resulting degeneracies in parameter fits (see Allen 2007), but they require assumptions about parametric functional forms. If completeness corrections are taken into account, it is important to ensure that different surveys use very similar or identical assumptions. In addition, because of the statistical nature of incompleteness corrections, only robust estimates of CF can be obtained since it cannot be determined around which primaries are the companions missed in the survey.

3 MULTIPLICITY ON THE MS

In this section, we review the multiplicity of Population I MS stars grouped by mass bins: solar-type stars ($M_{\star} \approx 0.7-1.3 \ M_{\odot}$; spectral types F through mid-K, Section 3.1), low-mass stars ($M_{\star} \approx 0.1-0.5 \ M_{\odot}$; M0–M6, Section 3.2), VLM stars ($M_{\star} \leq 0.1 \ M_{\odot}$; M8 and later, Section 3.3), intermediate-mass stars ($M_{\star} \approx 1.5-5 \ M_{\odot}$; B5–F2, Section 3.4) and high-mass stars ($M_{\star} \gtrsim 8 \ M_{\odot}$; B2 and earlier, Section 3.5). The mass dependence of stellar multiplicity is addressed in Section 5.1.

3.1 Solar-type stars

3.1.1 HISTORICAL OVERVIEW Solar-type stars represent an ideal population to probe stellar multiplicity on the MS. They are frequent enough to yield large, statistically robust samples and, while they are relatively bright and easy to observe, their luminosity contrast with any stellar companion is relatively modest. Solar-type stars were well represented in the samples of bright nearby stars studied over the last century (e.g., Heintz 1969), but completeness and sample homogeneity were poor. Even the large scale, uniform and dedicated solar-type multiplicity study conducted by Abt & Levy (1976) had a limited completeness and suffered from the biases inherent to its magnitude-limited nature.

The first modern, volume-limited multiplicity survey of solar-type stars was conducted by Duquennoy & Mayor (1991), who studied 164 objects out to 22 pc by collating results from many separate studies and obtaining their own new radial velocity measurements. For the last two decades, this study has represented the "gold standard" of multiplicity surveys among field stars, against which most subsequent multiplicity surveys compared their results. Nonetheless, significant incompleteness potentially affected the conclusions of Duquennoy & Mayor, prompting Raghavan et al. (2010) to revisit this topic with a volumelimited sample of 454 star out to 25 pc. In addition to a much cleaner sample, that survey built on much improved observational methods to reach a very high completeness rate ($\approx 97\%$). This study now represents the most complete multiplicity study of a well-defined sample and likely will remain a standard reference for years to come.

In addition to these global surveys, which consider companions at all orbital periods, valuable results have been obtained in large single-method surveys which enable more uniform detections rates. In the context of field solar-type stars, dedicated surveys for SBs, VBs and common proper motion companions have been conducted (e.g., Halbwachs et al. 2003, Mason et al. 1998, Tokovinin 2011).

3.1.2 MULTIPLICITY FREQUENCY While Abt & Levy (1976) concluded that "single stars are infrequent", both selection biases and survey incompleteness led these authors to significantly overestimate the companion frequency of solar-type stars. Instead of an average of 1.4 companion per primary star, Duquennoy & Mayor (1991) and Raghavan et al. (2010) demonstrated that $CF_{0.7-1.3 M_{\odot}}^{\rm MS} = 62 \pm 3\%$. In addition, Raghavan et al. estimated that $MF_{0.7-1.3 M_{\odot}}^{\rm MS} = 44 \pm 2\%$, i.e., a slight majority of all field solar-type stars are actually single, as is the Sun. Furthermore, Raghavan et al. took advantage of their large sample size to identify a dependency of multiplicity on stellar mass. Specifically, they found that that super-solar dwarfs have a marginally higher multiplicity rate than sub-solar dwarfs: $MF_{1-1.3 M_{\odot}}^{\rm MS} = 50 \pm 4\%$ vs $MF_{0.7-1 M_{\odot}}^{\rm MS} = 41 \pm 3\%$. Similarly, their sample leads to $CF_{1-1.3 M_{\odot}}^{\rm MS} = 75 \pm 5\%$ and $CF_{0.7-1 M_{\odot}}^{\rm MS} = 56 \pm 4\%$.

Mason et al. (1998) and Metchev & Hillenbrand (2009) have suggested that young ($\sim 0.1-1 \text{ Gyr}$) solar-type dwarfs have an excess of visual companions over their older counterparts. However, these surveys were skewed towards super-solar targets, reducing the amplitude of the proposed excess. Only a bias-free survey could conclusively demonstrate that young field solar-type stars host more stellar (and substellar) companions than their older MS counterparts.

3.1.3 ORBITAL PERIOD DISTRIBUTION The distribution of orbital periods for solartype dwarfs has long been known to be extremely broad, spanning up to 10 decades in orbital periods (Luyten 1930). While Öpik's law is a reasonable first approximation, the distribution cuts off at both extremes (Heintz 1969), although the exact location of the "breaks" are somewhat ill-defined (e.g., Poveda, Allen & Hernández-Alcántara 2007; Tokovinin & Lépine 2012). Indeed, the high completeness and large sample size of modern volume-limited samples now allows to confidently exclude Öpik's law as a good representation of the orbital period distribution over the entire range of periods. On the other hand, the log-normal description that was first proposed by Kuiper (1935b) has now become a clearly better description of the observed distribution (Raghavan et al. 2010). The maximum of the orbital period distribution occurs at $\overline{P} \approx 250$ yr ($\overline{a} \approx 45$ AU), with a dispersion of $\sigma_{\log P} \approx 2.3$.

3.1.4 MASS RATIO DISTRIBUTION The difficulties associated with the detection of lowmass companions has lead to widely discrepant conclusions regarding the overall mass ratio distribution of solar-type binaries. In turn, it has been proposed that this distribution is flat, bimodal, or monotonically increasing toward low-q systems (Trimble 1990, and references therein). Modern volume-limited surveys have clarified this ambiguous picture. Although Duquennoy & Mayor (1991) found a single peak around $q \approx 0.3$, the much higher completeness of the survey by Raghavan et al. (2010) firmly established that this distribution is instead flat down to $q \approx 0.1$ with the exception of a marginally significant peak at $q \gtrsim 0.95$. We fit a power law to the overall sample from Raghavan et al. (adding all three panels in their Figure 16) and obtain $\gamma_{0.7-1.3 M_{\odot}}^{\rm MS} = 0.28 \pm 0.05$.

At a more detailed level, it is apparent that the mass ratio distribution differs significantly between short- and long-period binaries. Splitting the sample of Raghavan et al. (2010) at the median orbital period, short-period binaries are characterized by a strong peak at $q \approx 1$ and a slowly declining f(q) function towards low mass ratios, while long-period binaries have a single peak around $q \approx 0.3$, similar to the total distribution found by Duquennoy & Mayor (1991). Specifically, we find that $\gamma_{0.7-1.3 M_{\odot}}^{\text{MS,log } P \leq 5.5} = 1.16 \pm 0.16$ and $\gamma_{0.7-1.3 M_{\odot}}^{\text{MS,log } P > 5.5} = -0.01 \pm 0.03$ (see also Tokovinin 2011), although neither power law is a perfect model of the observed distribution.

One remarkable feature of the mass ratio distribution for solar-type binaries is the dearth of substellar companions, the so-called "BD desert", first identified among SBs (Grether & Lineweaver 2006; Marcy, Cochran & Mayor 2000) and further proposed for VBs (e.g., Leconte et al. 2010). We discuss this in the general context of "extreme" mass ratio systems in Section 5.4.

3.1.5 HIGH-ORDER MULTIPLE SYSTEMS Triple and higher-order systems represent $\approx 25\%$ of all solar-type multiple systems, with a distribution of systems with *n* components that roughly follows a geometric distribution $N(n) \propto 2.5^{-n}$ up to n = 6 (see also Eggleton & Tokovinin 2008). However, this distribution is noticeably steeper for non-single systems: $N(n) \propto 3.7^{-n}$.

Not surprisingly, multiple systems are hierarchical. In all systems, the ratio of the orbital periods $P_{\rm long}/P_{\rm short}$ has a minimum value of $\gtrsim 5$ (e.g., Tokovinin 1997), consistent with long term stability arguments (Eggleton & Kiseleva 1995). The mass ratio distribution for the short period subsystems is flat up to $q \approx 0.9$, with a strong peak around $q \approx 1$ which is largely responsible for the marginal excess reported in the overall sample of Raghavan et al. (2010). On the other hand, the long period subsystems preferentially have $q \leq 0.5$ (Raghavan et al. 2010, Tokovinin 2008).

More than half of all pairs with $\log P \leq 2$ are members of high-order multiple systems and this proportion rises even further for the shortest-period systems (Allen et al. 2012, Tokovinin et al. 2006). In addition, the frequency of hierarchical triple systems among wide systems (separation \geq 1000 AU) is also high (Makarov, Zacharias & Hennessy 2008). This suggests that the presence of an outer companion plays an important role in the formation of tight pairs, presumably through energy and angular momentum exchanges. It is plausible that this phenomenon is also related to the formation of equal-mass subsystems.

3.2 Low-mass stars

3.2.1HISTORICAL OVERVIEW The high incidence of low-mass stars enables large samples of nearby objects to be searched for stellar companions. After the first attempts at detecting companions via the astrometric perturbation methods (e.g., Lippincott 1978), a series of instrumental breakthroughs over the last two decades have helped establish the multiplicity of low-mass stars through radial velocity (Marcy & Benitz 1989), direct imaging (Skrutskie, Forrest & Shure 1989) and speckle interferometry (Henry & McCarthy 1990) surveys. Although none of these surveys was volume-limited and their samples only partially overlaped, they allowed Fischer & Marcy (1992) to perform a complete analysis of M dwarf multiplicity for 166 stars that has long been used as the low-mass stars counterpart to the Duquennoy & Mayor (1991) survey for solar-type stars. More recently, similar-sized volume-limited surveys of low-mass stars have been conducted (Delfosse et al. 2004, Dieterich et al. 2012, Reid & Gizis 1997a). In particular, the RECONS consortium¹ has gathered a quasi-complete sample of stars and companions out to $\sim 10 \,\mathrm{pc}$ (Henry et al. 2006).

Multiplicity frequency While most studies agree that low-mass stars have 3.2.2fewer companions than solar-type stars, published multiplicity frequencies range from $26 \pm$ 9% (Leinert et al. 1997) to $42 \pm 9\%$ (Fischer & Marcy 1992). This spread can in part be explained by the small number of companions detected in these surveys, further compounded by significant incompleteness corrections in certain domains. Modern volume-limited surveys, with their near-perfect completeness, draw a much more coherent picture, where $MF_{0.1-0.5\ M_{\odot}}^{\rm MS} = 26 \pm 3\%$ and $CF_{0.1-0.5\ M_{\odot}}^{\rm MS} = 33 \pm 5\%$ (Delfosse et al. 2004, Dieterich et al. 2012, Reid & Gizis 1997a). The widely-used higher frequency reported by Fischer & Marcy (1992) can be traced back to their analysis of common proper motion binaries, which focused on the earliest spectral types (M3.5 and earlier). Indeed, recent studies have demonstrated that the frequency of wide companions $(50-10^5 \text{ AU})$ is a steep function of primary mass (Dhital et al. 2010) so that an average over all spectral types leads to a much reduced companion frequency at large separations. Despite this trend, there is no significant variation of the total multiplicity frequency between early-M and late-M dwarfs since most of their companions are found at separations shorter than 50 AU. On the other hand, the frequency of spectroscopic companions estimated by Fischer & Marcy has been upheld by more recent surveys (Clark, Blake & Knapp 2012)

3.2.3 ORBITAL PERIOD DISTRIBUTION Fischer & Marcy (1992) suggested that a lognormal distribution similar to that found for solar-type binaries could also apply to low-mass stars. As pointed out above, however, the frequency of wide companions was overestimated by Fischer & Marcy and the orbital period distribution must thus be revisited. A log-normal function is still acceptable for separation below 500 AU, with $\overline{a} \approx 5.3$ AU and $\sigma_{\log P} \approx$

¹http://www.recons.org/

1.3 based on the latest update to the RECONS sample. At larger separation, an Öpiklike distribution seems to hold out to 0.2–0.3 pc for early-M primaries (Dhital et al. 2010). Considering the small number of companions included in volume-limited samples, and the difficulty to draw a bias-free picture from the other available surveys, it remains unclear whether and how these two distributions are physically connected.

3.2.4MASS RATIO DISTRIBUTION The flat mass ratio distribution found by Fischer & Marcy (1992) suffered from potential biases as only primaries in the M0–M3 range and companions significantly above the substellar limit were considered. Nonetheless, including lower mass M-type primaries and all secondaries yields a mass ratio distribution that is flat or slightly declining towards low-q systems. For instance, we have fit a power law distribution to the total sample of Delfosse et al. (2004, their Figure 4) and found that $\gamma_{0.1-0.5 M_{\odot}}^{\text{MS}} = 0.39 \pm 0.23$. However, the situation is more subtle, since the mass ratio distribution is clearly dependent on the primary mass and the binary separation. Like solar-type binaries, short period M-dwarf binaries are significantly biased towards high-q systems (Reid & Gizis 1997a). In addition, low-q systems are limited to the earliest M-type dwarfs (Dieterich et al. 2012). Fitting the RECONS sample, we find that $\gamma_{0.1-0.5 M_{\odot}}^{\text{MS},a \le 5 AU} = 2.7 \pm 1.6$ and $\gamma_{0.1-0.5 M_{\odot}}^{\text{MS},a > 5 AU} = -0.3 \pm 0.3$. Similarly, we find that $\gamma_{0.3-0.5 M_{\odot}}^{\text{MS}} = -0.2 \pm 0.3$ and $\gamma_{0.1-0.3 M_{\odot}}^{\text{MS}} = 1.9 \pm 1.7$. Finally, we note that BD companions are easier to find around M dwarfs than around solar-type stars. Although dedicated searched have discovered a number of such companions (e.g., Burningham et al. 2009; Forveille et al. 2004; Nakajima, Oppenheimer & Kulkarni 1995; Radigan et al. 2008), the frequency of such companions is much lower than that of stellar companions based on volume-limited survey, with only 3 of the 23 companions in the RECONS sample being substellar.

3.2.5 HIGH-ORDER MULTIPLE SYSTEMS The ratio of multiple to binary systems among low-mass stars is $\approx 21\%$ (Reid & Hawley 2005), similar to the ratio observed for solar-type stars. Furthermore, the distribution of systems with n components among M dwarfs follows an $N(n) \propto 3.9^{-n}$ for $n \ge 2$, closely matching the distribution for solar-type systems². Finally, high-order multiple systems appear to have a much higher occurrence rate among systems which contain the widest pairs among M-dwarf systems (Law et al. 2010), suggesting that higher system masses are needed to produce the widest systems. Overall, besides the lower overall multiplicity of M dwarfs compared to solar-type stars, the basic statistics of high-order multiple systems do not seem to differ significantly between these two populations.

3.3 VLM stars and substellar objects

3.3.1 HISTORICAL OVERVIEW VLM objects, even the nearest ones, are typically very faint and therefore difficult targets for both high-resolution imaging and spectroscopic observations. Broad molecular bands and fast rotational velocities further hamper searches for SBs. In his review of BDs, Basri (2000) only briefly addressed the issue of multiplicity as the first BD binary systems had just been discovered. In the following decade, a slew of high-resolution surveys identified many systems, and thorough reviews of multiplicity in the substellar regime were presented by Burgasser et al. (2007) and Luhman (2012).

²See for instance the VLM binary database (http://www.vlmbinaries.org/)

3.3.2Multiplicity frequency In the absence of a large volume-limited survey for both VBs and SBs, a piece-wise approach is needed to estimate the total multiplicity frequency of VLM stars. The broad range of observed frequency of spectroscopic companions (Basri & Reiners 2006, Blake, Charbonneau & White 2010, Guenther & Wuchterl 2003, Reid et al. 2002, Tanner et al. 2012) is primarily a result of different completeness limits for each survey. Combining all these surveys, 77 objects of spectral type M8 or later have been observed at least three times with high-resolution spectroscopy, revealing 4 confirmed SBs and at least two more candidate binaries. The frequency of SBs ($a \leq 1 \,\mathrm{AU}$) is therefore at least 5.2^{+3.8}_{-1.5}%. At larger separations ($a \gtrsim 2 \,\text{AU}$), high-resolution imaging surveys indicate an observed binary frequency of about $15\pm3\%$ after correction for their inherent Malmquist bias (Bouy et al. 2003, Burgasser et al. 2007). It is likely that a handful of companions remain undetected in the 1–2.5 AU range (e.g., Bernat et al. 2010). Finally, no more than 2.3% of VLM objects have a companion in the 40-1000 AU range (Allen et al. 2007). Combining these various studies yields a total multiplicity fraction for field VLM objects of 20–25%, consistent with the frequency of $CF_{M_{\star} \leq 0.1 M_{\odot}}^{\text{field}} = 22^{+6}_{-4}\%$ estimated by Allen (2007) from a Bayesian analysis of several input surveys. We adopt this latter estimate, which takes into account survey incompleteness. Furthermore, while a few triple VLM systems have been confirmed or suspected (e.g., Artigau et al. 2011, Burgasser et al. 2012), too few of these have been discovered to draw a meaningful conclusion. Instead, we simply estimate that $CF_{M_{\star} \leq 0.1 M_{\odot}}^{\text{field}} \approx MF_{M_{\star} \leq 0.1 M_{\odot}}^{\text{field}}$. While the studies discussed so far generally focus on late-M or L-type objects, recent surveys found a similar multiplicity frequency for the coldest (T-type) BDs. Even the higher frequency of visual companions among "transition" objects (late-L to early-T dwarfs) can be explained by an enhanced Malmquist bias (Burgasser, Looper & Rayner 2010).

3.3.3 ORBITAL PERIOD DISTRIBUTION While too few VLM binaries are known to derive robust estimates, their orbital period distribution is much narrower than that of higher mass objects. Most companions are found with projected separations in the 1–10 AU range (see Burgasser et al. 2007). Only a handful of systems with projected separations larger than 50 AU are known (Billères et al. 2005, Burningham et al. 2010, Dhital et al. 2011) and this near-absence of wide systems cannot be accounted for by incompleteness (Allen 2007). Maxted & Jeffries (2005) and Allen (2007) have used statistical analyses to derive (truncated) log-normal analytical descriptions of the overall orbital period distribution. Both studies agree regarding the peak of the separation distribution ($\overline{a} \approx 4$ –7 AU), but differ on the width of the distribution. It is probable that an intermediate width ($\sigma_{\log P} \approx 0.4$ –0.5) is a good match to all observations. However, the orbital period distribution may follow a different analytical prescription and may even be asymmetric. More thorough searches for SBs are needed to settle this issue.

3.3.4 MASS RATIO DISTRIBUTION The mass ratio distribution for VLM stars³ is heavily skewed toward equal mass systems (see Burgasser et al. 2007). The vast majority of known systems have $q \ge 0.7$ and the most "extreme" systems have $q \approx 0.5$ (e.g., Biller et al. 2006, Kasper et al. 2007). Several studies have attempted to reproduce the observed mass ratio distribution by assuming an intrinsic power law distribution and taking into account selec-

³For substellar objects, estimated masses depend on the assumed age of the objects and are therefore not as precise as for hydrogen-burning stars. However, the uncertainty applies similarly to both components of a presumed coeval binary, so that absolute uncertainties in q are substantially smaller than on the masses of individual components.

tion and incompleteness effects (Allen 2007, Burgasser et al. 2006, Liu, Dupuy & Leggett 2010). While they all agree that $\gamma_{M_{\star} \leq 0.1 M_{\odot}} \leq 0$ can be firmly excluded, the estimated range of estimated values is very broad. This diversity is most likely the result of methodological differences. The shallow index found by Allen (2007) is unlikely to be appropriate as it leads to the prediction of many binaries with $q \leq 0.7$, which is at odds with deep imaging surveys. Instead, Burgasser et al. (2006) argues that the sharp decline in the observed distribution of mass ratios as q decreases is only modestly amplified by the Malmquist bias. While this issue is not entirely settled, we adopt the steep distribution proposed by Burgasser et al., $\gamma_{M_{\star} \leq 0.1 M_{\odot}} = 4.2 \pm 1.0$.

3.4 Intermediate-mass stars

3.4.1 HISTORICAL OVERVIEW Intermediate-mass stars are much more challenging for multiplicity studies than lower-mass stars. The steep mass-luminosity relationship makes low-mass companions hard to detect, and given typical distances, there is an un-probed gap between SBs and VBs. Further complicating the picture is the presence, among nearby field A dwarfs of some $\sim 30\%$ of chemically peculiar stars. Some of these categories owe their peculiarity to multiplicity (Abt 1965), inducing severe selection biases. In this review, we consider A stars as a unique population, averaging over the various sub-samples.

Searches for SBs among intermediate-mass stars have been ongoing for several decades (e.g., Abt & Snowden 1973, Carquillat & Prieur 2007, Carrier et al. 2002). Systematic searches for visual companions have been largely incomplete for a long time, although some of the studies listed above have attempted to include wider companions. The efforts to identify low-mass visual companions have accelerated in recent years as the presence of BD, and more recently planetary-mass, companions is investigated (e.g., Balega et al. 2011, Ivanov et al. 2006).

In parallel to these studies of field stars, another avenue to probe the multiplicity of intermediate-mass stars is to consider rich stellar associations in the process of dissipating in the field. By an age of 5 Myr, intermediate-mass stars already have reached the zero-age MS. The Scorpius-Centaurus OB association is the best studied region to date, both for SBs (e.g., Brown & Verschueren 1997) and visual companions (Kouwenhoven, Brown & Kaper 2007a, Shatsky & Tokovinin 2002), which allowed Kouwenhoven et al. (2007b) to perform a systematic analysis to derive the global multiplicity properties of intermediate-mass stars in that population.

3.4.2 MULTIPLICITY FREQUENCY The observed frequency of SBs among field intermediatemass stars lies in the 30–45% range (Abt 1983). In the Sco-Cen OB association, the fraction of SBs is at least 30%. The most thorough search for visual companions to A stars is the ongoing VAST survey (De Rosa et al. 2011). In the separation range 50–2000 AU, over which the survey is complete down to the substellar limit, they find a frequency of companions of $40\pm4\%$, although many of these companions have $q \leq 0.1$ and some of them may be unrelated background stars (R. De Rosa, priv. comm.). In the Sco-Cen OB association, Kouwenhoven et al. (2005) found a frequency of visual companions of 37%. Considering the limitations of current surveys, the Sco-Cen population provides the most robust estimate of the multiplicity of intermediate-mass stars. Current observations support the hypothesis that field stars have similar properties. Based on the various surveys listed above, we conclude that $MF_{1.5-5 M_{\odot}}^{\rm MS} \geq 50\%$, although incompleteness precludes robust estimates of the frequency of high-order multiplicity. In addition, the statistical analysis of Kouwenhoven et al. (2007b) resulted in $CF^{\rm MS}_{1.5-5\;M_{\odot}}=100\pm10\%.$

3.4.3 ORBITAL PERIOD DISTRIBUTION The orbital period distribution for field intermediatemass multiple systems extends from less than 1 d to projected separations of well beyond 10^4 AU (Abt 1983). Contrary to lower-mass field binaries, however, the orbital period distribution for A stars appears to be bimodal, with a peak among SBs around $P \sim 10$ d (e.g., Carquillat & Prieur 2007) and one for VBs around 350 AU (R. De Rosa, priv. comm.). In addition, there is marginal evidence for a dip around $P \sim 300$ d, with a tertiary peak or a broad plateau for separations in the 1–50 AU (Budaj 1999). Even though the observed distribution seems more complicated, Kouwenhoven et al. (2007b) tried both a power law and a log-normal representation for the period distribution of the Sco-Cen intermediatemass binaries with moderate success. In the former case, Öpik's law is favored, while in the latter formalism, they find $\sigma_{\log P} \gtrsim 2.3$. However, neither is a perfect description of the observed distribution and no simple analytical representation matches all observations.

3.4.4 MASS RATIO DISTRIBUTION The mass ratio distribution of field intermediatemass stars remains elusive as a result of incompleteness and selection biases, which are largely responsible for the early claims of a strong preference for high-q systems (e.g., Abt & Levy 1985). Taking those limitations into account, it now appears that the mass ratio distribution for SBs is either relatively shallow (e.g., $\gamma = -0.3 \pm 0.2$; Carquillat & Prieur 2007) or can be described by a broad Gaussian distribution centered around $q \approx 0.4$ (Carrier et al. 2002, Vuissoz & Debernardi 2004). Among visual companions, the VAST survey suggests a mass ratio distribution $\gamma \approx -0.6$ that is roughly consistent with that found for SBs (R. De Rosa, priv. comm.). Finally, we note that a similar mass ratio distribution was proposed by Shatsky & Tokovinin (2002) and Kouvenhoven et al. (2007b) for the Sco-Cen association. The latter study derived $\gamma_{1.5-5 M_{\odot}}^{\text{UpSco}} = -0.45 \pm 0.15$. Since this analysis includes both SBs and VBs, we adopt it as representative of all intermediate-mass multiple systems.

3.5 High-mass stars

3.5.1 HISTORICAL OVERVIEW The multiplicity of high-mass stars has long been the subject of intense scrutiny (Zinnecker & Yorke 2007, and references therein). However, their large distances from the Sun, large rotational velocity and the extremely high brightness contrast requirements conspire to draw an incomplete picture (Sana & Evans 2011). In addition, most high-mass stars are found in their birth environment, massive clusters or OB associations, and only $\sim 20\%$ of all nearby O stars are found in the field, many of which are thought to be ejected "runaway" stars (e.g., Chini et al. 2012, de Wit et al. 2004). As a result, studying only "field stars" is likely to yield a very biased picture.

High-mass SBs have been studied in all environments for several decades (Abt, Gomez & Levy 1990; Chini et al. 2012; Garmany, Conti & Massey 1980). Individual clusters and OB associations, despite smaller sample sizes, are more appropriate for a uniform characterization of multiple systems (Kiminki et al. 2012; Sana, Gosset & Evans 2009). Furthermore, it has recently become possible to probe the multiplicity of high-mass stars in the low-metallicity environment of the Large Magellanic Cloud (Sana et al. 2012b). Surveys for visual companions to high-mass stars have been conducted both for "all-sky" samples (e.g., Mason et al. 2009, Turner et al. 2008) and single clusters/associations (e.g., Duchêne et al. 2001, Peter et al. 2012).

The remarkably broad range of estimated frequency 3.5.2 Multiplicity frequency of spectroscopic companions among high-mass stars is due to a combination of small sample sizes, selection biases and incompleteness. Most recently, Sana et al. (2012a) estimated a bias- and completeness-corrected frequency of spectroscopic companions of $70\pm9\%$ out to $P \sim 3000 \,\mathrm{d}$ and down to $q \approx 0.1$. Early-B stars have a lower SB frequency than their higher-mass counterparts ($52 \pm 4\%$ for B2-B6 primaries; Chini et al. 2012). Due to the diversity of wavelengths and observing techniques, imaging surveys yield an even broader range of frequencies. A consensus has emerged that the frequency of visual companions over two decades in projected separations is $\sim 45 \pm 5\%$ (e.g., Peter et al. 2012, Turner et al. 2008). Incompleteness in these surveys is modest down to $q \approx 0.1$ overall but very uncertain below this limit. There is suggestive evidence that the frequency of visual companions decreases with the mass of primary (Peter et al. 2012, Preibisch et al. 1999). Interpolating between the populations of SBs and VBs suggests a "missed" companion frequency in the 10-20% range, in agreement with the small survey by Nelan et al. (2004). This leads to $CF_{M_{\star}\gtrsim 16}^{q\gtrsim 0.1} \approx 130 \pm 20\%$ and $CF_{8-16}^{q\gtrsim 0.1} \approx 100 \pm 20\%$. The latter is substantially lower than the estimate from Abt, Gomez & Levy (1990) but the large incompleteness correction tion of that survey probably explains this difference. In addition, the number of high-mass stars that are truly single stars is remarkably low. Considering only confirmed companions, $MF_{M_{\star} \gtrsim 16 M_{\odot}} \geq 80\%$ (Chini et al. 2012, Mason et al. 2009, Sana & Evans 2011). The situation is less well constrained for early-B stars, and $MF_{8-16 M_{\odot}} \ge 60\%$ (Abt, Gomez & Levy 1990) is a conservative lower limit since many tight visual companions were missed in that survey. High-mass stars in rich clusters and OB associations have essentially identical multiplicity frequencies, which in turn is significantly higher than that of field and runaway stars (Chini et al. 2012, Mason et al. 2009, Turner et al. 2008).

The relative lack of SBs with published orbital 3.5.3**Orbital Period Distribution** solutions has long impaired definitive analyses of the orbital period distribution for highmass multiple systems. The high frequency of short-period binaries proposed early on (Abt, Gomez & Levy 1990; Garmany, Conti & Massey 1980) was potentially subject to a strong selection bias, but has been confirmed by modern surveys, which find a pile-up of binaries with $4 d \le P \le 8 d$ (e.g., Sana et al. 2012a). Power law fits suggest a slowly-declining $(\alpha \approx -0.5)$ distribution out to P = 3000 d for both Galactic and Large Magellanic Cloud high-mass binaries (Sana et al. 2012a,b). However, extrapolating even a moderately steep distribution results in far too few visual companions, suggesting a more complex functional form. One possibility to match all observations combines two independent distributions: a population of short period binaries (log $P \lesssim 1$, 30% of all high-mass stars) and a power law period distribution extending out to $\gtrsim 10^4$ AU. While an Öpik-like law for the second population is compatible with available observations (Kiminki & Kobulnicky 2012), imaging surveys which repeatedly find a "peak" at the shortest probed angular separations rather support a slowly declining distribution instead (Mason et al. 2009, Nelan et al. 2004, Peter et al. 2012, Sana et al. 2011).

3.5.4 MASS RATIO DISTRIBUTION Limited sensitivity to even intermediate-mass companions has long plagued studies of the mass ratio distribution of high-mass stars. Furthermore, a large fraction of their companions are still in the PMS phase, resulting in model-dependent mass estimates. Both Kiminki & Kobulnicky (2012) and Sana et al. (2012a) concluded that the intrinsic distribution of mass ratios for high-mass SBs is essentially flat (e.g., $\gamma_{M_{\star}\gtrsim 16\,M_{\odot}}^{P\leq 3000\,d.q\geq 0.1} = -0.1\pm 0.6$ for Sana et al. 2012a, which we adopt here). While both studies

find a peak around $q \approx 0.8$, they rule out the presence of a separate population of equal-mass (so-called "twin") systems among SMC eclipsing binaries (Pinsonneault & Stanek 2006). Among VBs, Peter et al. (2012) concluded that a flat distribution and a broad Gaussian distribution centered on $q \sim 0.45$ match equally well their observations of high-mass stars in the Carina region. On the other hand, surveys of the Orion Nebula Cluster (ONC; Preibisch et al. 1999) and NGC 6611 (Duchêne et al. 2001) found a strong preference for low-mass companions ($q \leq 0.5$), although some of these systems may not be physically bound. Combining these three samples and fitting a power law to the resulting distribution yields $\gamma_{M_{\star}\gtrsim 16 M_{\odot}}^{a\geq 100 AU} = -0.55 \pm 0.13$, indeed favoring lower-mass companions but not significantly different from the distribution found for SBs down to $\sim 1 M_{\odot}$. Lower mass companions remain mostly out of reach of present surveys but the upcoming generation of high-contrast, high-resolution imaging instruments will soon provide a first foray into this category of "extreme" systems ($q \ll 0.1$).

4 MULTIPLICITY OF OLD AND YOUNG STARS

In this section, we consider multiplicity properties of stellar populations of old MS stars (Population II stars) and at increasing earlier stages of stellar evolution, from open clusters to the youngest embedded protostars. The time evolution of multiplicity is addressed in Section 5.2.

4.1 Population II stars

The multiplicity properties of Population II stars can be used to probe the star formation process in low metallicity environments. While early studies led to conflicting conclusions regarding a possible deficit of companions to Population II stars (Abt & Willmarth 1987, Jaschek & Jaschek 1959, Oort 1926), a more coherent picture has recently emerged thanks to instrumental improvements and larger and better-defined samples of low-metallicity dwarfs and high-velocity halo stars (e.g., Carney et al. 1994).

SBs among Population II stars appear to be equally frequent and to share very similar properties with their higher-metallicity counterparts (Goldberg et al. 2002, Latham et al. 2002). On the other hand, the frequency of VBs ($a \ge 10 \,\text{AU}$) is markedly lower than for solar-metallicity stars (Allen, Poveda & Herrera 2000; Lodieu et al. 2009; Zapatero Osorio & Martín 2004), though perhaps not for K-M dwarfs (Jao et al. 2009). As a result, the total multiplicity frequency of Population II stars is $CF_{0.7-1.3 \, M_{\odot}}^{Pop \, II} = 39 \pm 3 \,\%$ and $CF_{0.1-0.6 \, M_{\odot}}^{Pop \, II} = 26 \pm 6 \,\%$ (Jao et al. 2009, Rastegaev 2010, respectively).

The distribution of orbital period among solar-mass Population II multiple systems is characterized by a narrow peak in orbital period around log $P \approx 2-3$, accompanied by a longperiod tail out to ~ 10⁴ AU that follows roughly Öpik's law (Allen, Poveda & Herrera 2000) and a marked dearth of systems with orbital periods shorter than log $P \approx 1$ (Latham et al. 2002, Rastegaev 2010). The secondary peak at ~ 500 AU proposed by Zinnecker, Köhler & Jahreiß (2004) has not been confirmed by Rastegaev (2010), casting doubt on the its existence. Too few low-mass Population II binaries are known to robustly estimate their orbital period distribution, though they may represent a broader distribution than for solar-metallicity stars (Jao et al. 2009). Finally, while the mass ratio distribution of solar-type Population II binaries is still poorly determined, companion masses down to the substellar limit are found in roughly uniform distribution (Zapatero Osorio & Martín 2004).

4.2 Open clusters

Members of open clusters represent young (typically 50 Myr to 1 Gyr) and uniform populations that will eventually dissipate in the galactic field. These clusters are dynamically old, in the sense that their age is much larger than their crossing time, and wide binaries have long been disrupted (e.g., Kroupa 1995, Parker et al. 2009). Despite the advantage of large and homogeneous samples, multiplicity studies of open clusters are surprisingly scattered in the literature, with little overlap between samples. The best studied open clusters, in which both SBs and close VBs have been searched for, are the nearest ones ($D \leq 200 \text{ pc}$): α Per, the Pleiades, Praesepe and the Hyades. No dramatic difference has been identified between these clusters in terms of their multiplicity properties so that a composite average can be constructed to draw a global picture.

4.2.1 MULTIPLICITY FREQUENCY We first address the multiplicity of solar-type stars, which are the best-studied in open clusters. Most survey for SBs in open clusters agree on a 13–20% frequency of spectroscopic companions (e.g., Griffin 2012, Mermilliod, Grenon & Mayor 2008, Prosser 1992). Although adaptive optics surveys rely on substantial incompleteness correction for the closest companions ($\rho \leq 1''$), they are more sensitive to low-mass companions than speckle surveys (Mason et al. 1993a,b, Patience et al. 1998, 2002) and their companion frequency of ~ 25% over 1.5–2 decades in separation is more robust.

Combining SB and VB surveys therefore yields an observed companion frequency of open clusters of $\approx 40\%$ for solar-type stars, although some companions most likely still escape detection. Taking advantage of the range of distances to various clusters, Patience et al. (2002) found a total companion frequency of $48 \pm 5\%$ for all separations up to 580 AU and with mass ratios $q \gtrsim 0.25$. Assuming a mass ratio distribution similar to that of field FGK dwarfs ($\gamma \sim 0$), the total frequency is therefore $CF_{0.7-1.3 M_{\odot}}^{\text{open cluster}} \sim 65\%$. Independently, photometric studies have suggested multiplicity frequencies in the 65–70% range for solar-type stars (Converse & Stahler 2008, Kähler 1999).

Low-mass cluster members have only been systematically studied via spectroscopic and imaging surveys in the nearby Hyades cluster (Reid & Gizis 1997b, Reid & Mahoney 2000). Combining these two surveys leads to $CF_{0.1-0.5 M_{\odot}}^{\text{open cluster}} \approx 35 \pm 5\%$. Finally, VLM stars and BDs in open clusters have only been searched for visual companions (Bouy et al. 2006, Martín et al. 2003). Extrapolating the observed frequency of visual companions assuming an overall Gaussian orbital period distribution, Gizis & Reid (1995) suggested that $CF_{M_{\star} \lesssim 0.1 M_{\odot}}^{\text{open cluster}} = 27 \pm 16\%$. Photometric studies, which may be affected by a mass segregation selection effect, broadly support these results despite some significant scatter (Boudreault et al. 2012, Converse & Stahler 2008, Lodieu et al. 2007, Pinfield et al. 2003).

4.2.2 ORBITAL PERIOD DISTRIBUTION Although no single cluster has been studied over the entire range of orbital periods, it is possible to estimate the average orbital period distribution in open clusters in a piece-wise manner using the various surveys discussed above. The distribution for solar-type stars is broad and unimodal over the range $0 \leq \log P \leq 7$ (Patience et al. 2002). The second peak at $\log P \leq 1$ proposed by Griffin (2012) likely stems from the fact that their sample includes intermediate-mass stars.

The median separation of ~ 4 AU estimated by Patience et al. (2002) based on a lognormal fit, somewhat tighter than among field stars, may be a consequence of incompleteness among VBs. Indeed, the frequency of VBs included in their analysis by Patience et al. appears to be underestimated (Bouvier, Rigaut & Nadeau 1997; Bouvier et al. 2001; Morzinski et al. 2012), suggesting that the median separation is significantly larger. It is therefore probable that the orbital period distribution of solar-type binaries in open clusters is indistinguishable from that of field stars. Unfortunately, chance projection of unrelated members and background stars prevent any analysis of systems wider than $a \sim 1000$ AU in open clusters.

4.2.3 MASS RATIO DISTRIBUTION The distribution of mass ratios among solar-type binaries in open clusters is essentially flat for both SBs (e.g., Bender & Simon 2008, Goldberg & Mazeh 1994, Mermilliod & Mayor 1999) and VBs (Patience et al. 2002). In addition, we note that substellar companions to solar-type stars in open clusters are rare (Itoh et al. 2011, Metchev & Hillenbrand 2009), although a few have been recently discovered (Geißler et al. 2012, Rodriguez et al. 2012). On the other hand, multiple systems are increasingly skewed toward high-q for low-mass (Reid & Gizis 1997b, Reid & Mahoney 2000) and VLM (Bouy et al. 2006, Lodieu et al. 2007, Martín et al. 2003) members of open clusters.

4.3 Young nearby associations

The past decade has seen the discovery of several "moving groups", i.e., groups of comoving, coeval young stars that are unbound and not associated with a recognizable site of star formation (see Torres et al. 2008). The ages of these groups ($\sim 77-100$ Myr) make them a valuable bridge between the sites of ongoing star formation on one hand and open clusters and the field population on the other. Most of these groups are small $(N \leq 50)$, however, so they individually do not provide robust statistical constraints on the full binary population. Furthermore, many of the known binaries were discovered serendipitously rather than in statistically robust surveys. The few analyses to date suggest that they are broadly consistent with what is seen in the field and in young star-forming regions (e.g., Brandeker, Javawardhana & Najita 2003; Evans et al. 2012), except for an anomalously low frequency of wide binaries (a > 20 AU) in the η Cha moving group (Brandeker et al. 2006). Because of their youth and close proximity to the Sun ($d \leq 50$ pc), members of these associations have been subject to surveys aiming primarily at planet discovery (Chauvin et al. 2010, and references therein). These have revealed a small number of planetary-mass companions (Chauvin et al. 2004, Lagrange et al. 2009) and numerous BDs (e.g., Bowler et al. 2012, Lowrance et al. 1999, Neuhäuser & Guenther 2004, Wahhaj et al. 2011), as well as providing robust constraints on the existence and extent of the purported BD desert (Evans et al. 2012). A uniform analysis of all these results is still missing.

4.4 PMS stars

We now turn to multiplicity surveys targeting populations of PMS stars with ages in the 1–5 Myr range and with stellar mass $M_{\star} \leq 2 M_{\odot}$, collectively called T Tauri stars (TTS), or Class II/III sources depending on whether they host a protoplanetary disk (Lada 1987). Multiplicity surveys have targeted TTS located in a broad range of environments: dense stellar clusters (such as the ONC and IC 348), young OB associations (e.g., Sco-Cen), and "loose" (so-called T) associations (like the Taurus-Auriga or Chamaeleon star-forming regions). The latter populations typically contain many less stars (~ 100) than clusters and OB associations and only a handful of intermediate-mass stars. The closest of these populations are located ~ 150 pc away from the Sun.

Even though the existence of close pairs of TTS was HISTORICAL OVERVIEW 4.4.1noted very early with seeing-limited observations (Herbig 1962, Joy & van Briesbroeck 1944), the multiplicity of TTS was only addressed by dedicated surveys in the 1990s. Since then, thorough multi-wavelength surveys of nearby star-forming region have allowed the definition of large, complete samples for multiplicity studies (e.g., Herbig & Bell 1988, Luhman 2012). In addition to seeing-limited surveys (Reipurth & Zinnecker 1993) and mining of all-sky databases (Kraus & Hillenbrand 2009a), closer companions have been searched with a variety of high-angular resolution techniques, down to 2–3 AU resolution. In more or less chronological order, speckle interferometry (Ghez, Neugebauer & Matthews 1993: Leinert et al. 1993), lunar occultation (Simon et al. 1995), Hubble Space Telescope imaging (Padgett, Strom & Ghez 1997), adaptive optics (Duchêne, Bouvier & Simon 1999) and aperture masking (Kraus et al. 2008) have all been used in this context. The search for tighter companions remains challenging as strong and variable emission lines hamper radial velocity analyses. Nonetheless, a small number of SBs are known (e.g., Mathieu, Walter & Mvers 1989). The only systematic surveys published to date (Melo 2003, Nguyen et al. 2012) found a companion frequency similar to that of field stars but yielded no new orbital solutions. In the remainder of this section, we only discuss VB surveys, which have been much more extensively studied.

4.4.2 MULTIPLICITY FREQUENCY Binary surveys of PMS stars in T associations generally have measured companion frequencies that are roughly twice as high as among solar-type MS stars (e.g., Duchêne 1999). This led to the widespread meme that all young stars are born as multiples. The largest complete surveys to date have been conducted in Taurus and Upper Scorpius (Kraus et al. 2008, 2011). In Taurus, for instance, $CF_{0.7-2.5 M_{\odot}}^{\text{Class II/III,Tau}} = 64_{-9}^{+11}\%$ and $CF_{0.25-0.7 M_{\odot}}^{\text{Class II/III,Tau}} = 79_{-11}^{+12}\%$ over the separation range 3–5000 AU. Despite these high frequencies, it appears that ~ 1/4–1/3 of all systems consist of a single star, revealing the prevalence of high-order multiples. Kraus et al. (2011) noted that the distribution of degrees of multiplicity was broadly consistent with a purely Poissonian distribution, $N(n) \propto e^{-n}$, and hence that adding additional companions resembles a stochastic process. This distribution resembles the geometric distribution seen for the field (Section 3.1.5; Eggleton & Tokovinin 2008).

At the other extreme, numerous studies have shown that the multiplicity frequency in young dense clusters is much lower. This applies to the locally extreme ONC (Köhler et al. 2006, Petr et al. 1998), but also to other young clusters in the Orion complex (Beck, Simon & Close 2003) as well as to IC 348 in the Perseus cloud (Luhman, McLeod & Goldenson 2005). In the ONC, where the largest sample of targets has been assembled, Reipurth et al. (2007) found $CF^{\text{Class II/III,ONC}} = 8.8 \pm 1.1\%$ over a decade in projected separation, a factor of ~ 2.5 lower than found in Taurus over the same range. Although it has long been proposed that multiplicity is a smooth function of the stellar density, the statistical precision of current surveys in intermediate regions, such as the Ophiuchus cloud, do not conclusively support his scenario (King et al. 2012a, Marks & Kroupa 2012).

Although they long remained unaccessible, several surveys have targeted BDs and VLM stars in recent years (Biller et al. 2011, and references therein). Collectively, they demonstrated that most young low-mass objects have no companions at projected separations $\geq 5-10$ AU. Furthermore, Kraus & Hillenbrand (2012) performed a combined analysis of all published surveys and showed that the binary frequency declines precipitously from 0.5 M_{\odot} to below the substellar boundary, with $CF_{M_{\star} \lesssim 0.1\,M_{\odot}}^{\rm Class\,II/III} < 11\%$ (50% confidence)

4.4.3 ORBITAL PERIOD DISTRIBUTION As for the field population, both a log-normal and power law have been used with reasonable success to describe the distribution of orbital periods among TTS for the last two decades. The inability to distinguish between these distributions results in part from the limited range probed by observations. Recent studies suggest a separation distribution that follows Öpik's law out of ~ 5000 AU for solar-type stars and slowly declining for lower-mass stars (e.g., Kraus et al. 2011). As a result, the global distribution is consistent with both a very broad log-normal, with $\sigma_{\log P} \gtrsim 2$ (Kraus et al. 2011), or a slowly declining power law (e.g., King et al. 2012b). The latter conclusion also matches the results of Kouwenhoven et al. (2007b) for intermediate-mass multiple systems in Upper Scorpius. One important caveat is that the binary population in dense clusters drops sharply at projected separations of $\gtrsim 200$ AU (Reipurth et al. 2007; Scally, Clarke & McCaughrean 1999).

The orbital period distribution of binary TTS appears to be truncated, with very few systems outside of a maximum separation that varies smoothly with stellar mass (Kraus & Hillenbrand 2012, and references therein). The truncation radius appears to be ~ 300 AU for stars in the 0.25–0.5 M_{\odot} and as small as ~ 25 AU for the lowest mass TTS. However, the paucity is not complete: a small number of "unusually wide" pairs have been identified (Konopacky et al. 2007, Kraus et al. 2011, and references therein). These are discussed in a broader context in Section 5.5.

Finally, we note that the distribution of orbital separations is skewed towards tighter systems in disk-less Class III objects relative to Class II objects which host circumstellar disks. This is likely a result of disk evolution, since tidal forces in tight binaries ($a \leq 50 \text{ AU}$) can severely truncate disks and result in much shorter disk lifetime (Cieza et al. 2009, Kraus et al. 2012).

4.4.4 MASS RATIO DISTRIBUTION Given the shallow mass-luminosity relation for PMS stars, imaging surveys are quite sensitive to low-mass companions, routinely reaching $q \sim 0.1$ and more recently probing down to $q \leq 0.01$. However, uncertainties in PMS evolutionary models (Hillenbrand & White 2004) and the possibility of circumstellar dust emission and differential extinction could work to smooth out and/or bias the genuine mass ratio distributions, particularly for photometrically estimated mass ratios (Woitas, Leinert & Köhler 2001). Nonetheless, in the absence of complete spectroscopic follow-up campaigns (e.g., Hartigan & Kenyon 2003, White & Ghez 2001), this is the only available tool in most cases.

Most surveys of solar-type binary TTS have found mass ratio distributions that are roughly flat (Kraus et al. 2008, 2011, Lafrenière et al. 2008). Using a power law parametrization for the mass ratio distribution, Kraus et al. (2011) found $\gamma_{0.25-2.5 M_{\odot}}^{\text{Class II/III}} = 0.2 \pm 0.2$ for the entire Taurus population, albeit with a marginally significant trend with stellar mass, whereby lower mass systems preferentially have $q \gtrsim 0.5$. This trend is further expanded by considering the lowest mass binaries: Kraus & Hillenbrand (2012) found $\gamma_{0.07-0.12 M_{\odot}}^{\text{Class II/III}} \sim 1$. Due to the paucity of truly substellar binary systems, it is uncertain whether this trend continues through the substellar regime. We note, however, that there is a small fraction of objects that seem to host binary companions that are much less massive than the primary, and may also have unusually wide orbital radii (e.g., Luhman et al. 2009). These are further discussed in Section 5.5. At the other extreme, intermediate-mass binaries have a small but significant preference for low-q systems (Kouwenhoven et al. 2007b).

4.5 Embedded protostars

Over the last decade, increasing attention has been devoted to studying the multiplicity of embedded protostars, objects in the earliest phases of stellar evolution (e.g., Duchêne et al. 2007b). These represent the best chance of probing the formation mechanism of multiple systems at its origin, before subsequent internal and external dynamical evolution can affect unstable systems. However, protostars also pose serious challenges to companion searches. By definition, they are still embedded in the remnants of their nascent core and hence are best studied at infrared and radio wavelengths, where high-resolution techniques became available only recently. Furthermore, the mass of the central objects are extremely difficult to estimate.

Protostars are traditionally categorized based on their spectral energy distribution. Class 0 sources are the least evolved objects, with more mass in circumstellar material, mostly in the form of a dense envelope, than in the central object (André, Ward-Thompson & Barsony 1993). In contrast, the envelope mass of Class I sources is lower than that of the central object. We discuss both categories separately below.

4.5.1 CLASS I PROTOSTARS Although they are deeply embedded, Class I sources are bright near- and mid-infrared sources. To date, about 350 Class I sources have been observed with seeing-limited imaging (Connelley, Reipurth & Tokunaga 2008, Haisch et al. 2004) and, for about 100 of them, with high-resolution imaging techniques (Connelley, Reipurth & Tokunaga 2009, Duchêne et al. 2007a). Until a global statistical analysis of these various surveys is performed, we estimate the multiplicity of Class I in the following manner. First of all, we only consider sources within 500 pc of the Sun and companions with flux ratios $\Delta L \leq 4$ mag to limit incompleteness issues. The direct imaging survey of Connelley, Reipurth & Tokunaga (2008) yields 27 companions in the 200–2000 AU range out of 136 targets, while the adaptive optics surveys of Duchêne et al. (2007a) and Connelley, Reipurth & Tokunaga (2009) found 15 companions between 50 and 200 AU to 88 targets (with additional companions at shorter separations for the closest targets). Combining these two fractions, we find a total frequency of companions to Class I sources over the 50-2000 AU separation range of $CF^{\text{Class I}} = 36.9 \pm 5.2\%$. The associated distribution of projected separations is consistent with Öpik's law out to 5000 AU although chance projection of unrelated protostars could be non-negligible at the largest separations (Connelley, Reipurth & Tokunaga 2008).

There is marginal evidence for an environment dependency among Class I protobinaries in that the distribution of separations for Orion shows a marked peak in the 60–200 AU range not seen in other nearby star-forming regions (Connelley, Reipurth & Tokunaga 2009, Duchêne et al. 2007a). In this context, it is important to emphasize that Class I sources are distributed through the various molecular clouds in the Orion complex (a majority are in the L1641 cloud) and none belongs to the ONC. In other words, even the Orion Class I sources studied to date were born in dynamically quiet environments, so that any difference in that population is more likely to be a consequence of some other environmental difference than dynamical evolution. There is also evidence for a temporal evolution of the properties of protobinaries, as the most embedded, and presumably youngest, systems almost exclusively possess wide companions (1000 AU and wider), the frequency of which rapidly declines as one considers more evolved protostars (Connelley, Reipurth & Tokunaga 2008).

The search for much tighter companions among Class I protostars is remarkably challenging as radial velocity measurements are hampered by extremely weak photospheric features. Based on 2 or 3 measurements per systems (Viana Almeida et al. 2012) or comparing a protostar radial velocity to that of gas in its parent cloud (Covey et al. 2006), small-scaled surveys suggest that some systems tighter than a few AU have already formed by the Class I phase, but their interpretation remains ambiguous.

4.5.2 CLASS 0 PROTOSTARS The rarity of Class 0 sources makes it necessary to group all observed targets in a single sample to achieve meaningful statistics, precluding studies of environmental dependencies. In addition, high-resolution observations at near-infrared and centimeter wavelengths are probing scattered light and free-free emission (Reipurth 2000, Reipurth et al. 2004), respectively, neither of which can be used to unambiguously detect companions. Instead, the optimal observational probe of Class 0 multiplicity is therefore millimeter (broadly speaking, $850 \,\mu$ m-7 mm) interferometric mapping, even though the linear resolution typically achieved with such observations is only ~ 100 AU. Furthermore, clustering on scales of 1000 AU to 0.1 pc prevents studies of the widest systems (e.g., Carrasco-González et al. 2012; Peretto, Hennebelle & André 2007).

Early multiplicity studies of Class 0 protostars suggested that they have a very high multiplicity frequency although their target selection was mostly haphazard and potentially biased (e.g., Looney, Mundy & Welch 2000; Reipurth 2000). Recent high-resolution millimeter surveys, which primarily focused on the disk/envelope structure, reported low companion frequencies (Maury et al. 2010), albeit with extremely small sample size. Adding to the sample surveyed by Maury et al., Enoch et al. (2011) concluded that only 1 out of 11 targets had a companion within 2000 AU (albeit with minimum separations probed ranging from 50 to 400 AU). However, a much larger number of Class 0 sources have been imaged at high spatial resolution, leading to the discovery of several other companions in that separation range, thereby casting doubts on their apparent rarity (e.g., Looney, Mundy & Welch 2000). Further complicating the picture is the possibility that some continuum point sources could be related to outflow features instead of being bona fide companions (e.g., Maury, Ohashi & André 2012), a possibility that can only be clarified with follow-up observations. A global re-analysis of all high-resolution millimeter maps of Class 0 sources in the context of multiplicity is crucially missing but beyond the scope of this review. For now, we simply note that the multiplicity of Class 0 sources is not statistically significantly lower than that of the more evolved Class I protostars in the $\sim 100-2000$ AU range.

5 OVERALL TRENDS AND "EXTREME" SYSTEMS

We now gather the results discussed in the previous two sections to highlight the main trends associated with multiplicity as a function of primary mass and age. We then discuss in more detail several remarkable categories of companions: twin companions, extreme mass ratio systems and extremely wide systems. Possible physical implications of empirical trends are discussed in Section 6.

5.1 Multiplicity as a function of stellar mass

In this section, we focus on the mass-dependence of multiplicity properties on the MS (see Section 3 and Table 1). With continuously improving completeness and larger sample sizes, our empirical knowledge will soon allow decisive tests of the predictions from star formation theories.

Table 1: Multiplicity properties for Population I MS stars and field BDs			
Mass Range	Mult./Comp.	Mass Ratio	Orbital Period
	Frequency	Distribution	Distribution
$M_\star \lesssim 0.1M_\odot$	$MF = 22^{+6}_{-4} \%$	$\gamma = 4.2 \pm 1.0$	Unimodal (log-normal?)
	$CF = 22^{+6}_{-4} \%$		$\overline{a} \approx 4.5 \mathrm{AU}, \sigma_{\log P} \approx 0.5$
$0.1 M_{\odot} \lesssim M_{\star} \lesssim 0.5 M_{\odot}$	$MF = 26 \pm 3\%$	$\gamma = 0.4 \pm 0.2$	Unimodal (log-normal?)
	$CF=33\pm5\%$		$\overline{a} \approx 5.3 \mathrm{AU}, \sigma_{\log P} \approx 1.3$
$0.7 M_{\odot} \lesssim M_{\star} \lesssim 1.3 M_{\odot}$	$MF = 44 \pm 2\%$	$\gamma = 0.3 \pm 0.1$	Unimodal (log-normal)
	$CF=62\pm3\%$		$\overline{a} \approx 45 \mathrm{AU}, \sigma_{\log P} \approx 2.3$
$1.5 M_\odot \lesssim M_\star \lesssim 5 M_\odot$	$MF \ge 50\%$	$\gamma = -0.5 \pm 0.2$	Bimodal
	$CF = 100 \pm 10\%$		$\overline{P} \approx 10 \mathrm{d} \ \& \ \overline{a} \approx 350 \mathrm{AU}$
$8M_\odot \lesssim M_\star \lesssim 16M_\odot$	$MF \geq 60\%$		
	$CF = 100 \pm 20\%$		• • •
$M_{\star}\gtrsim 16M_{\odot}$	$MF \ge 80\%$	$\gamma^{P \le 3000 \mathrm{d}} = -0.1 \pm 0.6$	Peak + power law
	$CF = 130 \pm 20\%$	$\gamma^{a \ge 100 \mathrm{AU}} = -0.5 \pm 0.1$	$\overline{P} \approx 5 \mathrm{d} \& \mathrm{\ddot{O}pik} (?)$

5.1.1 MULTIPLICITY FREQUENCY As has long been known, the overall multiplicity frequency of MS stars is a steep, monotonic function of stellar mass (see Figure 1). Given the limited number of points available and the fact that companions to high-mass stars with $q \leq 0.1$ are not accounted for, we refrain from proposing a functional form that reproduces the intrinsic dependency of $CF(M_*)$. The dependency of $MF(M_*)$ is qualitatively similar to that of $CF(M_*)$ except for a shallower slope resulting primarily from the fact that MF has an intrinsic upper bound at 100%.

The frequency and order of multiple systems also are steep functions of stellar mass. For instance, all 11 sextuple systems in the Multiple Star Catalog are A-type stars (Eggleton & Tokovinin 2008), while a single such candidate system is known among nearby solar-type stars (Raghavan et al. 2010), and the highest-order multiple system containing only low-mass stars currently known has 4 components (Reid & Hawley 2005). While it appears that the ratio of binary to higher-order systems does not vary much for objects with $M_{\star} \leq 1.5 M_{\odot}$, it probably rises significantly toward high-mass stars.

5.1.2 ORBITAL PERIOD DISTRIBUTION The distribution of orbital periods is unimodal for solar-type and lower-mass stars, but both the median separation and width of the distribution decrease sharply with decreasing stellar mass. As a result, the frequency of companions in the 1–10 AU range (10–15%) does not vary significantly with stellar mass for $M_{\star} \leq 1.5 M_{\odot}$, including substellar objects. The much lower overall multiplicity of field BDs traces a marked deficit of wide companions as opposed to a uniform depletion over all separations.

Intermediate- and high-mass stars have more complex distributions of orbital periods. A strong peak at the shortest periods (log $P \approx 0$ –1) is found in both populations with an amplitude that increases with stellar mass. VBs show a peak for intermediate-mass stars, while the situation is less clear for high-mass stars, for which a shallow power-law is currently preferred. Interestingly, the frequency of companions in the 1–10 AU range among intermediate-mass appears in reasonable agreement with that observed among solar-type stars.



Figure 1:

Dependency of CF (red squares) and MF (blue triangles) with primary mass for MS stars and field VLM objects. The horizontal errorbars represent the approximate mass range for each population. For B and O stars, only companions down to $q \approx 0.1$ are included. The frequencies plotted here are the best-estimate numbers from Sections 3.1–3.5, also reported in Table 1.

5.1.3 MASS RATIO DISTRIBUTION Although a simple power law representation is imperfect for most samples, this formalism offers the most straightforward criterion to compare multiple systems of various masses. As shown in Figure 2, the observed distribution of mass ratios is close to a flat distribution ($|\gamma| \leq 0.5$) down to $q \sim 0.1$ for all masses $M_{\star} \geq 0.3 M_{\odot}$, extending to high-mass stars the conclusions of Reggiani & Meyer (2011). Below this limit, the mass ratio distribution becomes increasingly skewed towards high-q systems. Furthermore, shorter-period systems among solar-type and low-mass stars have a steeper mass ratio distribution than wider systems.

Many past studies have favorably compared the mass distribution of companions to that expected from random pairing from the IMF for field objects. In this situation, and leaving aside the "smearing" induced by broad ranges of primary masses (Tout 1991), γ should increase from -2.3 (Salpeter's slope) at the high-mass end to ~ 0 around or below the substellar limit, which does not match observations at any primary mass. Although a proper comparison between observations and theory/simulations should be conducted on the raw f(q) distributions rather than on estimated power law indices, the hypothesis of random pairing from the IMF is robustly excluded by observations (see Figure 2). On the other hand, the rarity of substellar companions to low-mass stars suggests an alternative model in which the mass ratio distribution is flat between q = 1 and a minimum companion mass, M_{sec}^{\min} , that is independent of the primary mass (at least up to $M_{\star} \sim 1.5 M_{\odot}$) and



Figure 2:

Power law index fitted to the observed distribution of mass ratios for multiple systems as a function of primary mass. Red diamonds represent fits to the overall population of multiple systems within a certain range of primary masses, whereas blue squares and green triangles represent fits to the subsets of "tight" $(P \leq \overline{P})$ and "wide" $(P \geq \overline{P})$ binaries, respectively. Horizontal errorbars represent the mass range for each subsample. The dashed curve represents the index that would be derived if the companions followed the single stars IMF of Chabrier (2003) and a simple power law would be fit to the resulting companion mass distribution over the mass ratio range $0.1 \leq q \leq 1$. The asterisk marks the derived power law index using the single stars IMF of Bochanski et al. (2010). The power law indices plotted here are the best-estimate numbers from Sections 3.1–3.5, also reported in Table 1.

lies around or somewhat below the substellar limit. However, given current statistical uncertainties, it is also possible that there is no such minimum companion mass, or that it varies with primary mass.

5.1.4 ECCENTRICITY DISTRIBUTION Early studies of the eccentricity distribution in stellar binaries were hampered by severe selection biases (e.g., Aitken 1932, Heintz 1969). More modern systematic surveys have now yielded a clear and uniform picture (e.g., Abt, Gomez & Levy 1990; De Rosa et al. 2012; Dupuy & Liu 2011; Kiminki & Kobulnicky 2012; Konopacky et al. 2010; Raghavan et al. 2010; Sana et al. 2012a).

The eccentricity distribution for field multiple systems shows remarkably little dependency on primary mass, extending the conclusions of Abt (2005, 2006) across the entire range of primary masses. Beyond a period of $\sim 100 \text{ d}$, all populations show an essentially flat



Figure 3:

Cumulative distribution of eccentricities for systems with $2 \le \log P \le 4$ for field multiple systems among solar-type stars (green curve; Raghavan et al. 2010), low-mass stars (orange curve; from the SB9 catalog, Pourbaix et al. 2004), VLM stars and BDs (red curve; Dupuy & Liu 2011), high-mass stars (blue curve; from the SB9 catalog and Abt 2005, Sana et al. 2012a). The dot-dashed curves indicate incomplete samples, for which the eccentricity distribution is potentially biased. The dashed and dotted curves represent the expected distribution for a flat and thermal distribution, f(e) = 2e, respectively.

distribution⁴. As illustrated in Figure 3, a broad Gaussian distribution centered on $\overline{e} \approx 0.4$ is a possible alternative representation, since it produces somewhat less high-eccentricity systems (e.g., Stepinski & Black 2001). Discriminating between these two representations hinges on large, unbiased samples, which are still missing for most stellar masses. In any case, the observed distributions are all inconsistent with the so-called "thermal distribution", even though the latter is frequently assumed in numerical simulations.

Almost all short-period systems have circular orbits as a result of tidal dissipation. The observed circularization periods, which range from $\leq 1 \text{ d}$ to $\approx 20 \text{ d}$, is a function of the primary mass and a monotonic function of stellar age, from open clusters members to Population II stars. A handful of exceptions with eccentric short-period binaries are known among solar-type binaries. All of them are either members of higher-order systems that may be undergoing Kozai cycles, and/or relatively young so that they have not fully circularized yet (Mathieu & Mazeh 1988).

Between these two regimes, the eccentricity distribution has a well-defined rising upper

⁴Although Duquennoy & Mayor (1991) favored such a distribution, it clearly violates their biascorrected distribution, which is flat beyond $e \approx 0.3$ (see their Fig. 6b).

envelope in the *P*-*e* plane. The mild correlation between mean eccentricity and orbital period, first proposed by Finsen (1936) and borne out by recent surveys, is mostly due to the dearth of systems with $e \gtrsim 0.6$ at orbital periods 10–300 day. Intriguingly, extrasolar planets populate the P - e plane in a very similar manner as stellar multiple systems (Udry & Santos 2007).

5.2 Multiplicity as a function of age

We now turn to the temporal evolution of multiplicity for solar-type and lower-mass stars, which we can probe for ages ranging from ≤ 0.1 Myr for the most embedded protostars to ≥ 10 Gyr for Population II field stars. Higher-mass stars are only probed over a much narrower age range and do not allow for as thorough an analysis.

5.2.1 MULTIPLICITY FREQUENCY No population other than nearby field stars has been sufficiently studied to draw a complete picture of multiplicity, so this analysis can only be achieved for certain categories of companions. For SBs, no significant difference is observed between TTS, open cluster members, Population I and II field stars, although this conclusion is potentially weakened by incompleteness and biases and deserves further attention. For the remainder of this section, "multiplicity frequency" specifically refers to the frequency of visual companions.

Even among VBs, only a narrow range of separations ($\sim 50-500$ AU) has been studied across all populations, due to the diversity of distances involved and observational techniques employed. This limited range is insufficient to draw statistically robust conclusions, given typical sample sizes. Instead, we proceed by estimating for each population the companion frequency per decade of projected separation using the widest possible separation range within the approximate 10–3000 AU range. This implicitly assume that all population of multiple systems follow Öpik's law. Although an imperfect fit to the overall observed distribution, it is a reasonable approximation in the separation range considered here (which is near the peak of most proposed log-normal separation distributions).

The resulting frequency of visual companions is shown in Figure 4, where we have elected to focus on the largest samples studied with uniform and complete techniques whenever possible. In addition, in cases where several surveys address the same "population" (e.g., several open clusters), we average the resulting multiplicity frequency and consider the dispersion of individual results as an additional source of uncertainty to account for some inherent diversity. Among TTS, we group separately dense clusters from loose associations, since their multiplicity frequency is markedly different.

Comparisons between populations of young stars and that of field stars is complicated by the broad range of stellar masses probed in the early stages of stellar evolution, where surveys typically include objects of all masses from the substellar limit up to $\sim 2 M_{\odot}$. In the past, it has been customary to compare most PMS multiplicity surveys to the wellstudied population of solar-type field stars. However, a fairer comparison should instead be performed with a weighted average of solar-type and low-mass stars that takes into account possible selection biases in the young stellar populations.

Figure 4 highlights two parallel trends. On the one hand, stellar multiplicity does not vary on timescales ≤ 50 Myr in low-density environment (star-forming regions and young associations). On the other hand, there is no significant difference in multiplicity frequency between young and open clusters on one hand, and Population I field star on the other hand. It remains undetermined whether the dichotomy of multiplicity properties at early ages nec-



Figure 4:

Dependency of the frequency of visual companions per decade of projected separation with age for solar-type (~ 0.7–1.5 M_{\odot} , blue triangles), low-mass (~ 0.1–0.5 M_{\odot} , red squares) and overall populations of young stars (~ 0.1–2 M_{\odot} , orange diamonds). Results from individual surveys are indicated as gray symbols and include Duchêne et al. (2007a) and Connelley, Reipurth & Tokunaga (2008) for Class I sources, Ratzka, Köhler & Leinert (2005), Kraus et al. (2008, 2011), and Lafrenière et al. (2008) for T/OB associations, Reipurth et al. (2007), Beck, Simon & Close (2003), Duchêne, Bouvier & Simon (1999) and Luhman, McLeod & Goldenson (2005) for clusters of TTS, Brandeker, Jayawardhana & Najita (2003); Brandeker et al. (2006), Chauvin et al. (2010) and McCarthy & White (2012) for nearby young associations, and Bouvier, Rigaut & Nadeau (1997); Bouvier et al. (2001) for open clusters. The horizontal errorbars represent the approximate age range for each population while vertical errorbars compound statistical uncertainties and the observed dispersion between similar regions. The dashed lines represent constant frequencies describing qualitatively the predicted behavior of low-density associations (top line) and dense clusters (bottom line). The hashed region represent the range of possible behaviors for stellar clusters in the embedded phases.

essarily implies that the initial multiplicity properties are environment-dependent. This is discussed in more detail in Section 6. Finally, the marked multiplicity deficit among Population II stars, which is limited to wide binaries, suggests either a metallicity dependence in binary formation or evolution on a timescale of several Gyr.

5.2.2 OTHER MULTIPLICITY PROPERTIES Beyond the evolution of the multiplicity frequency, most of the trends identified among PMS multiple systems closely match those observed among open cluster members and field stars. This includes the dependence of the mass ratio distribution on both stellar mass and separation, and the dependence of the max-

imum separation with stellar mass (e.g., Kraus et al. 2011). Possibly the most discrepant property when comparing young and mature multiple systems is the orbital period distribution: among young stellar objects it appears closer to Öpik's law than among field stars, especially for solar-type systems (e.g., Connelley, Reipurth & Tokunaga 2008; Kraus et al. 2011). However, this only applies for young multiple systems in loose associations. Given the marked deficit of wide companions in young clusters like the ONC, averaging the orbital period distribution over all star-forming regions (weighted by their relative star formation rates) would result in a distribution that is much more similar to that observed among field stars. Not surprisingly, it thus appears as if multiplicity properties of field stars can be explained from those of TTS without much evolution.

5.3 "Twin" binaries

The existence of a distinct population of "twin" binary systems has been repeatedly advocated and is still subject to intense debate, in part because of the loose definition of this phenomenon. Broadly speaking, twin binaries are near equal-mass, short-period binaries which are over-represented over the underlying population of binaries (e.g., Lucy & Ricco 1979).

An excess of systems with orbital period $\leq 43 \,\mathrm{d}$ (median period $\sim 7 \,\mathrm{d}$) and $q \geq 0.98$ among $0.5-2 \,M_{\odot}$ binaries has been detected (Lucy 2006, Simon & Obbie 2009) although this subpopulation represents only 2–3% of all SBs. This low frequency explains why it is not readily apparent in volume-limited surveys (Raghavan et al. 2010). It is probable that this excess results from accretion in the early stellar evolution stages that acts to equalize the components' masses.

Pinsonneault & Stanek (2006) found an excess of systems with $q \ge 0.87$ among eclipsing binaries among 7–27 M_{\odot} stars in the Small Magellanic Cloud. However, Lucy (2006) suggested that this trend could be explained in large part by selection biases and subsequent studies of Galactic high-mass stars do not find evidence for a separate population of twin systems (see Section 5.3). Still, it is plausible that twin binaries may be a feature associated with low metallicity, since high-mass binaries in the Large Magellanic Cloud show a marginally steeper mass ratio distribution than their Galactic counterpart (Sana et al. 2012b).

5.4 "Extreme" mass ratio systems

The interest in "extreme" mass ratio systems (broadly defined as $q \leq 0.1$) has rapidly grown, both in the context of the search for planetary-mass companions and because of the difficulty for binary formation model to produce them (e.g., Bate 2012). Recent surveys have largely overcome the contrast limitations of earlier searches, providing new insights into the formation of VLM companions. Figure 5, which is based on large-scale surveys of field and young stars, illustrates the marked deficit of low-q systems, especially among (very) low-mass objects.

The extremely low frequency of BD companions among solar-type SBs (Grether & Lineweaver 2006; Marcy, Cochran & Mayor 2000) is inconsistent with extrapolations of both the binary mass function and the planetary mass function (e.g., Howard et al. 2010, Reffert & Quirrenbach 2011, Sahlmann et al. 2011). This suggests a sharp discontinuity in companion masses around the hydrogen burning limit (Thies & Kroupa 2008), although this BD "desert" is not completely arid and warrants further analysis.



Figure 5:

Left: Distribution of mass ratios as function of mass for nearby field objects with $M_{\star} \leq 1.5 M_{\odot}$. The dotted and dot-dashed curves indicate constant companions masses at the substellar and planetary regime limits (0.075 and 0.013 M_{\odot}), respectively. Data for solar-type stars (green diamonds), low-mass stars (orange triangles) and VLM objects (red plus signs) are from Raghavan et al. (2010), the RECONS survey and Janson et al. (2012a), and the VLM binary database, respectively. BD companions to low-mass stars identified outside of large-scale surveys are shown as open orange triangles. The few systems with $M_{\text{prim}} \leq 0.5 M_{\odot}$ from Raghavan et al. (2010) correspond to lower mass subsystems in hierarchical multiple systems. Right: Similar plot for star-forming regions and young associations. Shown as gray diamonds in this plot are companions to Class II/III objects in Taurus, Upper Scorpius and Chamaeleon I, using the large surveys from Kraus et al. (2008, 2011), Kraus & Hillenbrand (2012), Lafrenière et al. (2008). In addition, red open diamonds represent a number of objects discovered via small-scale surveys or pointed observations with a particular emphasis on VLM primaries and/or companions.

Among visual companions, the stringent limits on the presence of substellar companions obtained a decade ago ($\leq 1\%$; e.g., Hinz et al. 2002, McCarthy & Zuckerman 2004, Oppenheimer et al. 2001) applied to samples largely dominated by low-mass stars. More recently, deeper surveys focusing on solar-type stars found potentially higher frequencies, although the very small number of detections limits the robustness of this result (Carson et al. 2006, Leconte et al. 2010, Metchev & Hillenbrand 2009, Tanner, Gelino & Law 2010). It remains unclear whether these substellar companions are a simple extension of the population of companions below $0.075 M_{\odot}$, as proposed by Metchev & Hillenbrand (2009), or whether a distinct trough in companion mass is present at large separations as for shortperiod systems. One approach to test whether there is a demarcating line between two distinct populations of visual companions consists in studying the frequency of (very) lowmass stellar companions to intermediate- and high-mass stars, which probe similar mass ratios. In this context, we note the existence of a handful of planetary-mass objects around intermediate-mass stars (Kalas et al. 2008, Marois et al. 2008) but an apparent dearth of companions around the substellar limit (Janson et al. 2011, Vigan et al. 2012). Upcoming surveys for wide planetary mass companions to nearby stars using "extreme" adaptive optics instruments will help address this issue in a statistically sound manner.

Multiple surveys have addressed the existence of the BD desert among visual companions in both star-forming regions and nearby associations (e.g., Evans et al. 2012; Kouwenhoven, Brown & Kaper 2007a; Tanner et al. 2007). These survey generally found that BD companions are rare but not absent (e.g., Janson et al. 2012b, Lafrenière et al. 2011, Nielsen et al. 2012). Finally, high-resolution imaging surveys also have discovered a small number of planetary-mass companions (~ 4–20 M_{Jup}) at distances of ≥ 100 AU from young stars in nearby star-forming regions (Chauvin et al. 2004, Ireland et al. 2011, Lafrenière, Jayawardhana & van Kerkwijk 2008, Luhman et al. 2006, Neuhäuser et al. 2005, Schmidt et al. 2008). Unfortunately, most of these discoveries were serendipitous, precluding a robust statistical analysis of their frequency.

5.5 Extremely wide systems

Observationally, the correlation between the median separation and primary mass observed for stars with $M_{\star} \leq 2 M_{\odot}$ is accompanied by a similar trend for the maximum binary separation a_{\max} (e.g., Reid et al. 2001). Although different functional forms and an unconfirmed discontinuity at ~ 0.2 M_{\odot} have been proposed (Burgasser et al. 2003, Close et al. 2003, Reid et al. 2001), this upper limit on binary separation appears to hold for both field populations and PMS systems (e.g., Faherty et al. 2011, Konopacky et al. 2007, Kraus & Hillenbrand 2009a). Furthermore, several authors used energetic arguments to propose that the maximum binary separation is actually set as a minimum binding energy (e.g. Close et al. 2003). It remains undetermined which quantity is the most physically relevant.

A key to understanding the physical meaning of the $a_{\rm max} - M_{\rm tot}$ correlation lies in "unusually wide" systems, which violate the general trend. While the vast majority of field binaries fall inside the maximum separation limit, numerous studies have successfully searched for systems well beyond that limit. For solar-type and intermediate-mass stars stars, systems with separation out to several pc have been identified (e.g., Caballero 2009, 2010, Shaya & Olling 2011). Among field low-mass stars, for which the term "unusually wide" was first coined, systems at separations $> 10^3$ AU have been found (e.g., Artigau et al. 2007, Caballero 2007, Dhital et al. 2010, Faherty et al. 2011, Radigan et al. 2009). A number of unusually wide low-mass systems have also been found in nearby star-forming regions (Allers et al. 2009, Béjar et al. 2008, Kraus & Hillenbrand 2009a, Luhman et al. 2009). Around the substellar limit, although a marked paucity of companions is observed beyond $\geq 25 \text{ AU}$, a small number of wide pairs have been identified, particularly among young objects (e.g., Béjar et al. 2008, Chauvin et al. 2004, Kraus & Hillenbrand 2007, Luhman et al. 2006). In all cases, these "unusually wide" systems have separations $a \gtrsim 10a_{\rm max}$. Even though follow-up observations of some of these wide systems indicate that they are frequently part of hierarchical multiple systems (Law et al. 2010), their binding energy is still well below the empirical limit discussed above.

The existence of these unusually wide systems, coupled with the fact that the empirical limit on binary separation has been repeatedly revised upwards over the last decade, suggests that there is no absolute upper bound on binary separation, but rather a gradual rarity beyond a certain separation.

6 IMPLICATIONS FOR BINARY FORMATION AND EVOLUTION

Multiplicity is a smooth function of stellar mass, both on the MS and in the earlier phases of stellar evolution. This trend points to a similar formation process across a broad range of prestellar core masses. Core ("prompt") fragmentation still appears as the leading candidate scenario to form multiple systems, although the extreme diversity of systems, most notably the large breadth of the orbital period distribution, may require at least two separate mechanisms (Tohline 2002).

Fragmentation appears to be mild, in that most cores only produce between 1 and 3 components. Since the frequency of quadruple and higher-order systems among field stars is low, most high-order systems would need to be destroyed early on if core fragmentation typically led to a larger number of stellar seeds. This destruction would result in a much lower overall multiplicity frequency among field stars, especially among low-mass stars, since many ejected components would be single stars. While this is supported by purely dynamical star formation simulations (e.g., Delgado-Donate et al. 2004), it is not borne out by observations (see also Goodwin & Kroupa 2005). Numerical simulations including the effects of magnetic field and/or radiative feedback appear to match more closely observed multiplicity properties since these processes severely hinder fragmentation (e.g., Bate 2012, Hennebelle & Teyssier 2008).

The statement that "most stellar systems formed in the Galaxy are likely single and not binary" (Lada 2006) is still undecided. While it is true that most field stars are single as a result of the preponderance of low-mass stars, it is contrary to observations in low-density star-forming regions. It is unclear whether these regions are representative of multiple star formation in all environments. Indeed, the universality of multiplicity properties among different birth environments is still in doubt. There are marked observational differences, most notably the lower multiplicity observed in clusters relative to loose associations of young stars. However, numerical simulations show that (internal and cluster) dynamical evolution is by and large over after 1 Myr (e.g., Marks & Kroupa 2012), suggesting that current observations do not probe multiplicity in an early enough evolutionary stage to be conclusive. Because frequent interactions in dense clusters can reduce the initial multiplicity frequency by a factor of 2–4, a picture in which multiplicity frequency is universally high at early times remains plausible (Kroupa & Petr-Gozens 2011). One critical test would be to determine the multiplicity properties of clustered protostars since Class I surveys have so far only targeted objects born in "quiet", low stellar density environments. Another possible test of this scenario consists in comparing the mass-dependency of multiplicity in young clusters and T/OB associations, since dynamical simulations predict that lower mass systems are more prone to disruption (Kroupa, Aarseth & Hurley 2001).

Dynamical evolution can naturally explain both the "low multiplicity" and "high multiplicity" tracks observed in Figure 4, in which clusters and loose associations represent extreme cases of highly effective and ineffective binary disruption (Kroupa & Bouvier 2003). Irrespective of whether initial multiplicity properties are universal, the existence of these two tracks has immediate implications on the origin of field stars. Indeed, it appears that most stars form in clusters, an argument already made in the past based on multiplicity (Marks & Kroupa 2011, Parker et al. 2009, Patience et al. 2002) as well as from star/cluster counts (Lada & Lada 2003). A more thorough analysis that goes beyond the multiplicity frequency and also considers other parameters, such as the orbital period distribution or dependencies on the primary mass, is still needed.

The fact that the mass ratio distribution is essentially flat for most primary masses

 $(M_{\star} \gtrsim 0.3 M_{\odot})$ but becomes increasingly steeper at lower masses (Section 5.1.3) sets important constraints on both binary formation itself and the subsequent accretion through which final stellar masses are built. In particular, the suggestion that there may be a lower limit to companion masses, which remains unconfirmed with current observations, could be pointing to the smallest scale associated with fragmentation. In addition, the (small but growing) population of planetary-mass objects at large distances from their parent star (Section 5.4) poses a significant challenge to existing models of star and planet formation. Their orbital radii are too large for a "planetary" formation model (i.e., from a protoplanetary disk; Boss 2001, Pollack et al. 1996), but their masses are extremely low for a "binary" formation model (falling near or below the opacity-limited minimum mass; Kratter Murray-Clay & Youdin2010)). Nonetheless, it appears that one of these unlikely outcomes must occur. Determining whether these objects represent an extension of (sub)stellar companions to extremely low masses or form a separate population of objects will be addressed in the next few years.

At short orbital periods, although the long-discussed BD desert has been repeatedly confirmed as tracing physically distinct formation mechanisms, the remarkable similitude between the eccentricity distributions of planetary and stellar-like companions (including massive BDs) is intriguing. One possible interpretation is that the final eccentricity in both planetary and multiple stellar systems is ultimately set by dynamical *N*-body interactions. If this is true, it remains to be understood why the eccentricity distribution of multiple systems does not follow the thermal distribution, however.

The observed correlation between primary mass and "maximum" binary separation is plausibly linked to the typical prestellar core size (both on the order of ~ 0.1 pc Bahcall & Soneira 1981, Lada et al. 2008). Similarly, the median binary separation, which also scales with primary mass, could be hinting at a preferred spatial scale for fragmentation. Alternatively, the physical origin of a_{max} could lie in the dynamical evolution of systems. Indeed, the widest systems have a very weak binding energy that makes them vulnerable to disruption by other nearby passing stars, either in their birth environment (Kroupa 1995, Parker et al. 2009) or in the Galactic field (Jiang & Tremain 2010; Weinberg, Shapiro & Wasserman 1987). The latter secular evolution may be responsible for the lower multiplicity observed among Population II stars at large separations. A constant minimum binding energy can be expected in dynamically relaxed populations, such as virialized stellar clusters or field stars. However, even dynamically primordial environments like Taurus-Auriga and Upper Scorpius broadly follow the same a_{max} trend, casting doubt on this interpretation.

The physical origin of unusually wide systems (Section 5.5) is not fully ascertained. They could form in a similar manner as other binaries from a single core. Since they would not survive dynamical interactions in dense clusters, one possible explanation for their extreme paucity in the field population is that "unusually wide" binaries form via a low-probability channel, and that only those binary systems born in unbound associations can remain bound until they reach the field (Kraus et al. 2008, Kraus & Hillenbrand 2009a). The main $M_{tot} - a_{max}$ relation would then indicate the typical constraint from protostellar core sizes or the maximum spatial scale for fragmentation. Alternatively, they may have formed from separate cores and have been ejected simultaneously from a cluster on sufficiently similar trajectories as to remain weakly bound (Kouwenhoven et al. 2010, Moeckel & Clarke 2011). The low probability of this scenario readily accounts for their rarity among field stars.

7 SUMMARY POINTS

- The frequency and key properties associated with stellar multiplicity vary smoothly with primary mass, suggesting a similar formation process across all core masses.
- Multiplicity properties appear to be mostly set at the PMS phase. Subsequent evolution is modest at best, as expected from dynamical evolution models.
- No single population of binary systems has a mass-ratio distribution that agrees with random pairing from the IMF. Also, no population has an eccentricity distribution that follows the "thermal" distribution, f(e) = 2e.
- Observations of both PMS and field stars do not support the existence of a firm upper limit on binary separation, but an increasing rarefaction towards the largest scales (e.g., ~ 1 pc for solar-type systems).
- At short orbital periods (log $P \leq 3$), there is a marked distinction between planetaryand stellar-type systems, presumably tracing different formation mechanisms.

8 FUTURE ISSUES

- Are the multiplicity properties resulting from the star formation process universal or do they depend on the native environment? The mass-dependency of multiplicity among PMS in different environments stars holds a critical clue.
- Are there significant differences in the multiplicity properties of high-mass stars relative to lower-mass stars that would not follow a smooth mass dependency? A significant frequency of stellar companions with $q \leq 0.1$, or the confirmation of a separate but substantial population of "twin" binaries, would set high-mass stars apart.
- What is the formation path and associated timescale for SBs? The frequency and properties of VBs appear to be set by the TTS phase, but is this also true for SBs?
- What is the origin and fate of "unusually wide" systems?
- Is the dearth of "extreme mass ratio" systems (known as the BD desert among tight solar-type and lower-mass binaries) independent of separation and primary mass? What are the physical processes responsible for this phenomenon?
- Does multiplicity hinder or promote planet formation? How does it depend on binary separation, mass ratio, and eccentricity?

9 LITERATURE CITED

LITERATURE CITED

- Abt HA 1965. Ap. J. Suppl. 11:429
- Abt HA, Snowden MS 1973. Ap. J. Suppl. 25:137
- Abt HA, Levy SG 1976. Ap. J. Suppl. 30:273
- Abt HA 1983. Annu. Rev. Astron. Astrophys. 21:343
- Abt HA, Levy SG 1985. Ap. J. Suppl. 59:229
- Abt HA, Willmarth DW 1987. Ap. J. 318:786
- Abt HA, Gomez AE, Levy SG 1990. Ap. J. Suppl. 74:551
- Abt HA 2005. Ap. J. 629:507
- Abt HA 2006. Ap. J. 651:1151
- Aitken RG 1932. MNRAS 92:596
- Allen C, Poveda A, Herrera MA 2000. Astron. Astrophys. 32:385
- Allen PR, Koerner DW, McElwain MW, Cruz KL, Reid IN 2007. AJ 133:971
- Allen PR 2007. Ap. J. 668:492
- Allen PR, Burgasser AJ, Faherty JK, Kirkpatrick JD 2012. AJ 144:62
- Allers KN, Liu MC, Shkolnik E, Cushing MC, Dupuy TJ, Mathews GS et al. 2009. Ap. J. 697:824
- Ambartsumian V 1937. Astr. Zh. 14:207
- André P, Ward-Thompson D, Barsony M 1993. Ap. J. 406:122
- André P, Ward-Thompson D, Barsony M 2000. In *Protostars and Planets IV*, ed. V Mannings, AP Boss, SS Russell, p. 59. Tucson: University of Arizona Press
- Artigau E, Lafrenière D, Doyon R, Albert L, Nadeau N, Robert J 2007. Ap. J. Lett. 659:L49
- Artigau E, Lafrenière D, Doyon R, Liu M, Dupuy TJ et al. 2011. Ap. J. 739:48
- Bahcall JN, Soneira RM 1981. Ap. J. 246:122
- Balega YY, Dyachenko VV, Maksimov AF, Mologolovets EV, Rastegaev DA, Romanyuk II 2011. Astron. Nach. 332:978
- Basri G 2000. Annu. Rev. Astron. Astrophys. 38:485
- Basri G, Reiners A 2006. Astron. J. 132:663
- Bate MR 2012. MNRAS 419:3115
- Beck TL, Simon M, Close LM 2003. Ap. J. 583:358
- Béjar VJS, Zapatero Osorio MR, Pérez-Garrido A, Martín EL, Rebolo R et al. 2008. Ap. J. Lett. 673:L185
- Bender CF, Simon M 2008. Ap. J. 689:416
- Bernat D, Bouchez AH, Ireland M, Tuthill P, Martinache F et al. 2010. Ap. J. 715:724
- Billères M, Delfosse X, Beuzit J-L, Forveille T, Marchal L, Martín EL 2005. Astron. Astrophys. Lett. 440:L55
- Biller BA, Kasper M, Close LM, Brandner W, Kellner S 2006. Ap. J. Lett. 641:L141
- Biller BA, Allers K, Liu MC, Close LM, Dupuy T 2011. Ap. J. 740:39
- Blake CH, Charbonneau D, White RJ 2010. Ap. J. 723:684
- Bochanski JJ, Hawley SL, Covey KR, West AA, Reid IN et al. 2010. Astron. J. 139:2679 Boss AP 2001. Ap. J. 563:367
- Boudreault S, Lodieu N, Deacon NR, Hambly NC 2012. MNRAS 426:3419
- Bouvier J, Rigaut F, Nadeau D 1997. Astron. Astrophys. 323:139
- Bouvier J, Duchêne G, Mermilliod J-C, Simon T 2001. Astron. Astrophys. 375:989
- Bouy H, Brandner W, Martín EL, Delfosse X, Allard F, Basri G 2003. Astron. J. 126:1526
- Bouy H, Moraux E, Bouvier J, Brandner W, Martín EL et al. 2006. Ap. J. 637:1056
- Bowler BP, Liu MC, Shkolnik EL, Tamura M 2012. Ap. J. 756:69
- Brandeker A, Jayawardhana R, Najita J 2003. Astron. J. 126:2009
- Brandeker A, Jayawardhana R, Khavari P, Haisch KE, Mardones D 2006. Ap. J. 652:1572
- Brown AGA, Verschueren W 1997. Astron. Astrophys. 319:811
- Budaj J, 1999. MNRAS 310:419
- Burgasser AJ, Kirkpatrick JD, Reid IN, Brown ME, Miskey CL, Gizis JE 2003. Ap. J. 586:512
- Burgasser AJ, Kirkpartick JD, Cruz KL, Reid IN, Leggett SK et al. 2006. Ap. J. Suppl. 166:585
- Burgasser AJ, Reid IN, Siegler N, Close L, Allen P et al. 2007. In Protostars and Planets V, ed. B

Reipurth, D Jewitt, K Keil, p. 427. Tucson: University of Arizona Press Burgasser AJ, Looper D, Rayner JT 2010. Astron. J. 139:2448 Burgasser AJ, Luk C, Dhital S, Bardalez GD, Nicholls CP et al. 2012. Ap. J. 757:110 Burningham B, Pinfield DJ, Leggett SK, Tinney CG, Liu MC et al. 2009. MNRAS 395:1237 Burningham B, Leggett SK, Lucas PW, Pinfield DJ, Smart RL et al. 2010. MNRAS 404:1952 Caballero JA 2007. Astron. Astrophys. Lett. 462:L61 Caballero JA 2009. Astron. Astrophys. 507:251 Caballero JA 2010. Astron. Astrophys. 514:98 Carney BW, Latham DW, Laird JB, Aguilar LA 1994. Astron. J. 107:2240 Carquillat J-M, Prieur J-L 2007. MNRAS 380:1064 Carrasco-González C, Osorio M, Anglada G, D'Alessio P., Rodríguez LF et al. 2012. Ap. J. 746:71 Carrier F, North P, Udry S, Babel J 2002. Astron. Astrophys. 394:151 Carson JC, Eikenberry SS, Smith JJ, Cordes JM 2006. Astron. J. 132:1146 Chabrier G 2003. Publ. Astron. Soc. Pac. 115:763 Chauvin G, Lagrange A-M, Dumas C, Zuckerman B, Mouillet D et al. 2004. Astron. Astrophys. Lett. 425:L29 Chauvin G, Lagrange A-M, Bonavita M, Zuckerman B, Dumas C et al. 2010. Astron. Astrophys. 509:52Chini R, Hoffmeister VH, Nasseri A, Stahl A, Zinnecker H 2012. MNRAS 424:1925 Cieza LA, Padgett DL, Allen LE, McCabe CE, Brooke TY et al. 2009. Ap. J.l 696:L84 Clark BM, Blake CH, Knapp GR 2012. Ap. J. 744:119 Close LM, Siegler N, Freed M, Biller B 2003. Ap. J. 587:407 Connelley MS, Reipurth B, Tokunaga A 2008. Astron. J. 135:2526 Connelley MS, Reipurth B, Tokunaga A 2009. Astron. J. 138:1193 Converse JM, Stahler SW 2008. Ap. J. 678:431 Covey KR, Greene TP, Doppmann GW, Lada CJ 2006. Astron. J. 131:512 Delfosse X, Beuzit J-L, Marchal L, Bonfils X, Perrier C et al. 2004. In ASP Conf. Series 318:166 Delgado-Donate EJ, Clarke CJ, Bate MR, Hodgkin ST 2004. MNRAS 351:617 De Rosa RJ, Bulger J, Patience J, Leland B, Macintosh B et al. 2011. MNRAS 415:854 De Rosa RJ, Patience J, Vigan A, Wilson PA, Schneider A et al. 2012. MNRAS 422:2765 de Wit WJ, Testi L, Palla F, Vanzi L, Zinnecker H 2004. Astron. Astrophys. 425:937 Dhital S, West AA, Stassun KG, Bochanski JJ 2010. Astron. J. 139:2566 Dhital S, Burgasser AJ, Looper DL, Stassun KG 2011. Astron. J. 141:7 Dieterich SB, Henry TJ, Golimowski DA, Krist JE, Tanner AM 2012. Astron. J. 144:64 Duchêne G, Bouvier J, Simon T 1999. Astron. Astrophys. 343:831 Duchêne G 1999. Astron. Astrophys. 341:547 Duchêne G, Simon T, Eislöffel J, Bouvier J 2001. Astron. Astrophys. 379:147 Duchêne G, Bontemps S, Bouvier J, ANdré P, Djupvik AA, Ghez AM 2007a. Astron. Astrophys. 476:229 Duchêne G, Delgado-Donate E, Haisch KE, Loinard L, Rodríguez LF 2007b. In Protostars and Planets V, ed. B Reipurth, D Jewitt, K Keil, p. 379. Tucson: University of Arizona Press Duquennoy A, Mayor M 1991. Astron. Astrophys. 248:485 Dupuy TJ, Liu MC 2011. Ap. J. 733:122 Eggleton PP, Kiseleva L 1995. Ap. J. 455:640 Eggleton PP, Tokovinin AA 2008. MNRAS 389:869 Enoch ML, Corder S, Duchêne G, Bock DC, Bolatto AD et al. 2011. Ap. J. Suppl. 195:21 Evans TM, Ireland MJ, Kraus AL, Martinache F, Stewart P et al. 2012. Ap. J. 744:120 Faherty JK, Burgasser AJ, Bochanski JJ, Looper DL, West AA et al. 2011. Astron. J. 141:71 Finsen WS 1936. MNRAS 96:862 Fischer DA, Marcy GW 1992. Ap. J. 396:178 Forveille T, Ségransan D, Delorme P, Martín EL, Delfosse X et al. 2004. Astron. Astrophys. Lett. 427:L1

- Fröbrich D 2005. Ap. J. Suppl. 156:169
- Garmany CD, Conti PS, Massey P 1980. Ap. J. 242:1063
- Geißler K, Metchev SA, Pham A, Larkin JE, McElwain M, Hillenbrand LA 2012. Ap. J. 746:44
- Gerbaldi M, Floquet M, Hauck B 1985. Astron. Astrophys. 146:341
- Ghez AM, Neugebauer G, Matthews K 1993. Astron. J. 106:2005
- Gizis JE, Reid IN 1995. Astron. J. 110:1248
- Goldberg D, Mazeh T 1994. Astron. Astrophys. 282:801
- Goldberg D, Mazeh T, Latham DW, Stefanik RP, Carney BW, Laird JB 2002. Astron. J. 124:1132
- Goodwin SP, Kroupa P 2005. Astron. Astrophys. 439:565
- Grether D, Lineweaver CH 2006. Ap. J. 640:1051
- Griffin RF 2012. JAA 33:29
- Guenther EW, Wuchterl G 2003. Astron. Astrophys. 401:677
- Haisch KE, Greene TP, Barsony M, Stahler SW 2004. Astron. J. 127:1747
- Halbwachs J-L, Mayor M, Udry S, Arenou F 2003. Astron. Astrophys. 159:175
- Hale A 1994. Astron. J. 107:306
- Hartigan P, Kenyon SJ 2003. Ap. J. 583:334
- Heacox WD 1996. Publ. Astron. Soc. Pac. 108:591
- Heintz WD 1969. JRASC 63:275
- Hennebelle P, Teyssier R 2008. Astron. Astrophys. 477:25
- Henry TJ, McCarthy DW 1990. Ap. J. 350:334
- Henry TJ, Jao W-C, Subasavage JP Beaulieu TD, Ianna PA et al. 2006. Astron. J. 132:2360
- Herbig GH 1962. Adv. Astr. Astr. 1:47
- Herbig GW, Bell KR 1988. Lick Obs. Bull. 1111
- Hillenbrand LA, White RJ 2004. Ap. J. 604:741
- Hinz JL, McCarthy DW, Simons DA, Henry TJ, Kirkpatrick JD, McGuire PC 2002. Astron. J. 123:2032
- Howard AW, Marcy GW, Johnson JA, Fischer DA, Wright JT et al. 2010. Science 330:653
- Ireland MJ, Kraus A, Martinache F, Law N, Hillenbrad LA 2011. Ap. J. 726:113
- Itoh Y, Oasa Y, Funayama H, Hayashi M, Fukagawa M et al. 2011. Research in Astron. Astrophys. 11:335
- Ivanov VD, Chauvin G, Foellmi C, Hartung M, Huélamo N et al. 2011. Astrophys. Space Sci. 304:247
- Janson M, Bonavita M, Klahr H, Lafrenière D, Jayawardhana R, Zinnecker H 2011. Ap. J. 736:89
- Janson M, Hormuth F, Bergfors C, Brandner W, Hippler S et al. 2012a. Ap. J. 754:44
- Janson M, Jayardhana R, Girard JH, Lafrenière D, Bonavita M et al. 2012b. Ap. J. Lett. 758:L2
- Jao W-C, Mason BD, Hartkopf WI, Henry TJ, Ramos SN 2009. Astron. J. 137:3800
- Jaschek C, Jaschek M 1959. Zeitschrift Astrophys. 48:263
- Jiang Y-F, Tremaine S 2010. MNRAS 401:977
- Joy AH, van Briesbroeck G 1944. Publ. Astron. Soc. Pac. 56:123
- Kähler H 1999. Astron. Astrophys. 346:67
- Kalas P, Graham JR, Chiang E, Fitzgerald MP, Clampin M et al. 2008. Science 322:1345
- Kasper M, Apai D, Janson M, Brandner W 2007. Astron. Astrophys. 472:321
- Kiminki DC, Kobulnicky HA, Ewing I, Bagley Kiminki MM, Lundquist M et al. 2012. Ap. J. 747:41 Kiminki DC, Kobulnicky HA 2012. Ap. J. 751:4
- King RR, Parker RJ, Patience J, Goodwin SP 2012a. MNRAS 421:2025
- King RR, Goodwin SP, Parker RJ, Patience J 2012b. MNRAS in press (arxiv:1209.2415)
- Koch RH, Hrivnak BJ 1981. Astron. J. 86:438
- Köhler R, Petr-Gozens MG, McCaughrean MJ, Bouvier J, Duchêne G et al. 2006. Astron. Astrophys. 458:461
- Konopacky QM, Ghez AM, Rice EL, Duchêne G 2007. Ap. J. 663:394
- Konopacky QM, Ghez AM, Barman TS, Rice EL, Bailey JI et al. 2010. Ap. J. 711:1087
- Kouwenhoven MBN, Brown AGA, Zinnecker H, Kaper L, Portegies Zwart SF 2005. Astron. Astro-

phys. 137:154

Kouwenhoven MBN, Brown AGA, Kaper L 2007a. Astron. Astrophys. 464:581

- Kouwenhoven MBN, Brown AGA, Portegies Zwart SF, Kaper L 2007b. Astron. Astrophys. 474:77
- Kouwenhoven MBN, Goodwin SP, Parker RP, Davies MB, Malmberg D, Kroupa P 2010. MNRAS

404:1835

Kratter KM, Murray-Clay RA, Youdin AN 2010. Ap. J. 710:1375

Kraus AL, Hillenbrand LA 2007. Ap. J. 664:1167

Kraus AL, Ireland MJ, Martinache F, Lloyd JP 2008. Ap. J. 679:762

Kraus AL, Hillenbrand LA 2009. Ap. J. 703:1511

Kraus AL, Ireland MJ, Martinache F, Hillenbrand LA 2011. Ap. J. 731:8

Kraus AL, Hillenbrand LA 2012. Ap. J. 757:141

Kraus AL, Ireland MJ, Hillenbrand LA, Martinache F 2012. Ap. J. 745:9

Kroupa P 1995. MNRAS 277:1491

Kroupa P, Aarseth S, Hurley J 2001. MNRAS 321:699

Kroupa P, Bouvier J 2003. MNRAS 346:343

Kroupa P, Petr-Gozens MG 2011. Astron. Astrophys. 529:92

Kuiper GP 1935a. Publ. Astron. Soc. Pac. 47:15

- Kuiper GP 1935b. Publ. Astron. Soc. Pac. 47:121
- Lada CJ 1987. In *Star forming region Proc. IAU Symp. 115*, ed M Peimber J Jugaku, p. 1, Dordrecht: D Reidel Publishing

Lada CJ, Lada EA 2003. Annu. Rev. Astron. Astrophys. 41:57

Lada CJ 2006. Ap. J. 640:L63

Lada CJ, Muench AA, Rathborne J, Alves JF, Lombardi M 2008. Ap. J. 672:410

Lafrenière D, Jayawardhana R, Brandeker A, Ahmic M, van Kerkwijk MH 2008. Ap. J. 683:844

Lafrenière D, Jayawardhana R, van Kerkwijk MH 2008. Ap. J. Lett. 689:L153

Lafrenière D, Jayawardhana R, Janson M, Helling C, Witte S, Hauschildt P 2011. Ap. J. 730:42

Lagrange A-M, Gratadour D, Chauvin G, Fusco T, Ehrenreich D et al. 2009. Astron. Astrophys. Lett. 493:L21

Latham DW, Stefanik RP, Torres G, David RJ, Mazeh T et al. 2002. Astron. J. 124:1144

Law NM, Dhital S, Kraus A, Stassun KG, West AA 2010. Ap. J. 720:1727

Leconte J, Soummer R, Hinkley S, Oppenheimer BR, Sivaramakrishnan A et al. 2010. Ap. J. 716:1551

Leinert C, Zinnecker H, Weitzel N, Christou J, Ridgway ST et al. 1993. Astron. Astrophys. 278:129

Leinert C, Henry T, Glindermann A, McCarthy DW, 1997. Astron. Astrophys. 325:159

Lippincott SL 1978. Space Science Rev. 22:153

Liu MC, Dupuy TJ, Leggett SK 2010. Ap. J. 722:311

Lodieu N, Dobbie PD, Deacon NR, Hodgkin ST, Hambly NC, Jameson RF 2007. MNRAS 380:712

Lodieu N, Zapatero Osorio MR, Martín EL 2009. Astron. Astrophys. 499:729

Looney LW, Mundy L, Welch WJ 2000. Ap. J. 529:477

Lowrance PJ, McCarthy C, Becklin EE, Zuckerman B, Schneider G et al. 1999. Ap. J. Lett. 512:L69

Lucy LB, Ricco E 1979. Astron. J. 84:401

Lucy LB 2006. Astron. Astrophys. 629:635

Luhman KL, McLeod KK, Goldenson N 2005. Ap. J. 623:1141

Luhman KL, Wilson JC, Brandner W, Skrutskie MF, Nelson MJ et al. 2006. $Ap.\ J.$ 649:894

Luhman KL, Mamajek EE, Allen PR, Muench AA, Finkbeiner DP 2009. Ap. J. 691:1265

Luhman KL 2012. Annu. Rev. Astron. Astrophys. 50:65

Luyten WJ 1930. Pub. Nat. Academy Science 16:257

Maís Appellániz J 2010. Astron. Astrophys. 518:1

Makarov VV, Zacharias N, Hennessy GS 2008. Ap. J. 687:566

Marcy GW, Benitz KJ 1989. Ap. J. 344:441

Marcy GW, Cochran WD, Mayor M 2000. In *Protostars and Planets IV*, ed. V Mannings, AP Boss, SS Russell, p. 1285. Tucson: University of Arizona Press

- Marks M, Kroupa P 2011. MNRAS 417:1702
- Marks M, Kroupa P 2012. Astron. Astrophys. 543:8
- Marois C, Macintosh B, Bartman T, Zuckerman B, Song I et al. 2008. Science 322:1348
- Martín EL, Barrado y Navascués D, Baraffe I, Bouy H, Dahm S 2003. Ap. J. 594:525
- Mason BD, McAlister HA, Hartkopf WI, Bagnuolo WG 1993a. Astron. J. 105:220
- Mason BD, Hartkopf WI, McAlister HA, Sowell JR 1993b, Astron. J. 106:637
- Mason BD, Henry TJ, Hartkopf WI, Ten Brumelaar T, Soderblom DR 1998. Astron. J. 116:2975
- Mason BD, Hartkopf WI, Giez DR, Henry TJ, Helsel JW 2009. Astron. J. 137:3358
- Mathieu RD, Mazeh T 1988. Ap. J. 326:256
- Mathieu RD, Walter FM, Myers PC 1989. Astron. J. 98:987
- Mathieu RD 1994. Annu. Rev. Astron. Astrophys. 32:465
- Maury A, André P, Hennebelle P, Motte F, Stamatellos D et al. 2010. Astron. Astrophys. 512:40
- Maury A, Ohashi N, André P 2012. Astron. Astrophys. 539:130
- Maxted PFL, Jeffries RD 2005. MNRAS Lett. 362:L45
- McCarthy C, Zuckerman B 2004. Astron. J. 127:2871
- McCarthy K, White RJ 2012. Astron. J. 143:134
- Melo CHF 2003. Astron. Astrophys. 410:269
- Mermilliod J-C, Mayor M 1999. Astron. Astrophys. 352:479
- Mermilliod J-C, Grenon M, Mayor M 2008. Astron. Astrophys. 491:951
- Metchev SA, Hillenbrand LA 2009. Ap. J. Suppl. 181:62
- Moeckel N, Clarke CJ 2011. MNRAS 415:1179
- Morzinski KM, Macintosh BA, Close LM, Marois C, Konopacky Q, patience J 2012. Adaptive Optics Systems III – SPIE Conf. Series, 8447:160
- Nakajima T, Oppenheimer BR, Kulkarni SR 1995. Nature 378:463
- Nelan EP, Walborn NR, Wallace DJ, Moffat AFJ, Makidon RB et al. 2004. Astron. J. 128:323
- Neuhäuser R, Guenther EW 2004. Astron. Astrophys. 420:647
- Neuhäuser R, Guenther E, Wuchterl G, Mugrauer M, Bedalov A, Hauschildt PH 2005. Astron. Astrophys. Lett. 435:L13
- Nguyen DC, Brandeker A, van Kerkwijk MH, Jayawardhana R, 2012. Ap. J. 745:119
- Nielsen EL, Liu MC, Wahhaj Z, Biller BA, Hayward TL et al. 2012. Ap. J. 750:53
- Oort JH 1926. Publ. Kaypten Astron. Lab. Gröningen 40:1
- Oppenheimer BR, Golimowski DA, Kulkarni SR, Matthews K, Nakajima T et al. 2001. Astron. J. 121:2189
- Öpik E 1924. Publ. Obs. Tartu 25:6
- Padgett DL, Strom SE, Ghez AM 1997. Ap. J. 477:705
- Parker RJ, Goodwin SP, Kroupa P, Kouwenhoven MBN 2009. MNRAS 397:1577
- Patience J, Ghez AM, Reid IN, Weinberger AJ, Matthews K 1998. Astron. J. 115:1972
- Patience J, Ghez AM, Reid IN, Matthews K 2002. Astron. J. 123:1570
- Peretto N, Hennebelle P, André P 2007. Astron. Astrophys. 464:983
- Peter D, Feldt M, Henning T, Hormuth F 2012. Astron. Astrophys. 538:74
- Petr MG, Coudé du Foresto V, Beckwith SVW, Richichi A, McCaughrean MJ 1998. Ap. J. 500:825
- Pinfield DJ, Dobbie PD, Jameson RF, Steele IA, Jones HRA, Katsiyannis AC 2003. MNRAS 342:1241
- Pinsonneault MH, Stanek KZ 2006. Ap. J. Lett. 639:L67
- Pollack JB, Hubickyj O, Bodenheimer P, Lissauer JJ, Podolak M, Greenzweig Y 1996. Icarus 124:62
- Popovic GM, Pavlovic R, Ninkovic S 2004. Astron. J. 169:59
- Pourbaix D, Tokovinin AA, Batten AH, Fekel FC, Hartkopf WI et al. 2004. Astron. Astrophys. 424:727
- Poveda A, Allen C, Hernández-Alcántara A 2007. Binary stars as critical tools & tests in contemporary astrophysics – Proc. IAU Symp. 240, ed WI Hartkopf, EF Guinan, P Harmanec, p. 417, Cambridge: Cambridge University Press
- Preibisch T, Balega Y, Hofmann K-H, Weigelt G, Zinnecker H 1999. New Astron. 4:531

- Prosser CF 1992. Astron. J. 103:488
- Radigan J, Lafrenière D, Jayawardhana R, Doyon R 2008. Ap. J. 689:471
- Radigan J, Lafrenière D, Jayawardhana R, Doyon R 2009. Ap. J. 698:405
- Raghavan D, McAlister HA, Henry TJ, Latham DW, Marcy GW et al. 2010. Ap. J. Suppl. 190:1
- Rastegaev DA 2010. Astron. J. 140:2013
- Ratzka T, Köhler R, Leinert C 2005. Astron. Astrophys. 437:611
- Reffert S, Quirrenbach A 2011. Astron. Astrophys. 527:140
- Reggiani MM, Meyer MR 2011. Ap. J. 738:60
- Reid IN, Gizis JE 1997a. Astron. J. 113:2246
- Reid IN, Gizis JE 1997b. Astron. J. 114:1992
- Reid IN, Mahoney S 2000. MNRAS 316:827
- Reid IN, Gizis JE, Kirkpatrick JD, Koerner DW 2001. Astron. J. 121:489
- Reid IN, Kirkpatrick JD, Liebert J, Gizis JE, Dahn CC, Monet DG 2002. Astron. J. 124:519
- Reid IN, Hawley SL 2005. New light on dark stars Red dwarfs, low-mass stars, brown dwarfs, Chichester: Praxis Publishing, 594pp, 2nd ed.
- Reipurth B, Zinnecker H 1993. Astron. Astrophys. 278:81
- Reipurth B 2000. Astron. J. 120:3177
- Reipurth B, Rodíguez LF, Anglada G, Bally J 2004. Astron. J. 127:1736
- Reipurth B, Guimarães MM, Connelley MS, Bally J 2007. Astron. J. 134:2272
- Rodriguez DR, Marois C, Zuckerman B, Macintosh B, Melis C 2012. Ap. J. 748:30
- Sahlmann J, Ségransan D, Queloz D, Udry S, Santos NC et al. 2011. Astron. Astrophys. 525:95
- Sana H, Gosset E, Evans CJ 2009. MNRAS 400:1479
- Sana H, Evans CJ 2011., In Active OB stars: structure, evolution, mass loss and critical limits Proc. IAU Symp. 272, ed C Neiner, G Wade, G Meynet, G Peters, p. 474, San Francico: Astron. Soc. Pac.
- Sana H, Lacour S, Le Bouquin J-B, de Koter A, Moni Bidin C et al. 2011. In Four decades of research on massive stars – ASP Conf. Series, ed L Drissen, C Robert, N St-Louis, AFJ Moffat, in press (arXiv:1109.6654)
- Sana H, de Mink SE, de Koter A, Langer N, Evans CJ et al. 2012a. Science 337:444
- Sana H, de Koter A, de Mink SE, Dunstall PR, Evans CJ et al. 2012b. Astron. Astrophys., in press (arxiv:1209.4638)
- Scally A, Clarke C, McCaughrean MJ 1999. MNRAS 306:253
- Schmidt TOB, Neuhäuser R, Seifhart A, Vogt N, Bedalov A et al. 2008. Astron. Astrophys. 491:311
- Shatsky N, Tokovinin A 2010. Astron. Astrophys. 382:92
- Shaya EJ, Olling RP 2011. $Ap.\ J.S$ 192:2
- Simon M, Ghez AM, Leinert C, Cassar L, Chen WP et al. 1995. Ap. J. 443:625
- Simon M, Obbie RC 2009. Astron. J. 137:3442
- Skrustkie MF, Forrest WJ, Shure M 1989. Astron. J. 98:1409
- Stepinski TF, Black DC 2001. Astron. Astrophys. 371:250
- Sterzik MF, Tokovinin AA 2002. Astron. Astrophys. 384:1030
- Tanner A, Beichman C, Akeson R, Ghez A, Grankin KN et al. 2007. Publ. Astron. Soc. Pac. 119:747
- Tanner AM, Gelino CR, Law NM 2010. Publ. Astron. Soc. Pac. 122:1195
- Tanner AM, White R, Bailey J, Blake C, Blake G et al. 2012. Ap. J.S 203:10
- Thies I, Kroupa P 2008. MNRAS 390:1200
- Tohline JE 2002. Annu. Rev. Astron. Astrophys. 40:349
- Tokovinin AA 1997. Astron. Astrophys. 124:75
- Tokovinin A, Thomas S, Sterzik M, Udry S 2006. Astron. Astrophys. 450:681
- Tokovinin A 2008. *MNRAS* 389:925
- Tokovinin A 2011. Astron. J. 141:52
- Tokovinin A, Lépine S 2012. Astron. J. 144:102
- Torres CAO, Quast GR, Melo CHF, Sterzik MF 2008. In *Handbook of star formation*, ed B Reipurth, p. 757, San Francico: Astron. Soc. Pac.

Tout CA 1991. MNRAS 250:701

- Trimble V 1990. MNRAS 242:79
- Turner NH, ten Brummelaar TA, Roberts LC, Mason BD, Hartkopf WI, Gies DR 2008. Astron. J. 136:554
- Udry S, Santos NC 2007. Annu. Rev. Astron. Astrophys. 45:397
- van de Kamp P 1975. Annu. Rev. Astron. Astrophys. 13:295
- Viana Almeida P, Melo C, Santos NC, Figueira P, Sterzik M, Gameiro JF 2012. Astron. Astrophys. 539:62
- Vigan A, Patience J, Marois C, Bonavita M, De Rosa, RJ et al. 2012. Astron. Astrophys. 544:9
- Vuissoz C, Debernardi Y 2004. In Spectroscopically and spatially resolveing the components of close binary stars – ASP Conf. Series 318, ed RW Hilditch, H Hensberge, K Pavlovski, p. 320, San Francisco: Astron. Soc. Pac.
- Wahhaj Z, Liu MC, Biller BA, Clarke F, Nielsen EL et al 2011. Ap. J. 729:139
- Weinberg MD, Shapiro SL, Wasserman I 1987. Ap. J. 312:367

White RJ, Ghez AM 2001. Ap. J. 556:265

- Woitas J, Leinert C, Köhler R 2001. Astron. Astrophys. 376:982
- Zapatero Osorio MR, Martín EL 2004. Astron. Astrophys. 419:167
- Zinnecker H, Köhler R, Jahreiß H 2004. In The environment and evolution of double and multiple stars – IAU Colloq. 191, ed C Allen, C Scarfe, Rev. Mex. Astron. Astrofís. 21:33
- Zinnecker H, Yorke HW 2007. Annu. Rev. Astron. Astrophys. 45:481

10 SIDEBAR

Note to editor: place alongside Section 2

Short period binaries (log $P \lesssim 4$) are best identified through radial velocity monitoring. Longer period ("visual") systems are revealed by direct and increasingly higher-angular resolution imaging surveys using speckle interferometry, adaptive optics-aided imaging, the Hubble Space Telescope, aperture masking on monolithic telescopes, lunar occultation and long-baseline interferometry. Companions in the most challenging "intermediate" regime $(P \gtrsim 5 \text{ yr}, \rho \lesssim 0.05")$ can also be probed by astrometric monitoring, although it has been difficult to apply to large, well-defined stellar populations except for bright stars observed by *Hipparcos*.

Multiplicity can also be probed indirectly. In uniform stellar populations, high mass ratio systems appear over-luminous, i.e., located up to 0.75 mag above the sequence of single objects. The vertical spread in the HR diagram can thus be modeled statistically to constrain the multiplicity frequency and the mass ratio distribution. Alternatively, individual objects whose spectrum contains two distinct sets of photospheric features, revealing the presence of a companion of much different effective temperature, are known as "spectral" binaries. These are easiest to detect in cases where the $M_{\star} - T_{\rm eff}$ relationship is steep but the $M_{\star} - L_{\star}$ is shallow, most notably among VLM stars and BDs.

11 Acronyms and Definition

- a: semi-major axis (expressed in AU unless otherwise stated)
- BD: brown dwarf
- CF: companion frequency (average number of companion per target in a population)
- IMF: initial mass function
- MF: multiplicity frequency (fraction of multiple systems in a population)
- MS: Main Sequence
- ONC: Orion Nebula Cluster
- P: orbital period (expressed in days unless otherwise stated)
- PMS: Pre-Main Sequence
- SB: spectroscopic binary (broadly speaking, $\log P \lesssim 3$)
- TTS: T Tauri stars
- VB: visual binary (broadly speaking, $\rho \gtrsim 0.05$ ")
- VLM: very low-mass
- α : index of the power-law fit to the orbital period distribution
- γ : index of the power-law fit to the mass ratio distribution
- ρ : angular separation of a visual binary (expressed in arcsec unless otherwise stated)