Occultation Newsletter

Volume VI, Number 1

November, 1993

ISBN 0737-6766

Published quarterly by the International Occultation Timing Association Joan Bixby Dunham, Editor

Occultation Newsletter

Volume VI, Number 1

November, 1993

ISBN 0737-6766

Occultation Newsletter is published by the International Occultation Timing Association. Editor: Joan Bixby Dunham; 7006 Megan Lane; Greenbelt, MD 20770-3012; U.S.A. Please send editorial matters to the above. Send new and renewal memberships and subscriptions, back issue requests, address changes, graze prediction requests, reimbursement requests, special requests, and other IOTA business, but not observation reports, to: Craig and Terri McManus; 2760 SW Jewell Ave.; Topeka, KS 66611-1614; U.S.A.

FROM THE PUBLISHER

For subscription purposes, this is the third issuee of 1993. It is the first issue of Volume 6. IOTA annual membership dues, including ON and supplements for U.S.A., Canada, and Mexico \$25.00 for all others 30.00

Annual IOTA membership dues may be paid by check drawn on an American bank, money order, cash, or by charge to Visa or MasterCard. If you use Visa or MasterCard, include your account number, the expiration date, and your signature.

ON subscription (1 year = 4 issues)

for U.S.A., Canada, and Mexico	20.00
for all others	25.00

Single issues are 1/4 of the price shown.

Although they are available to IOTA members without charge, nonmembers must pay for these items:

Local circumstance (asteroidal appulse) predictions 1.00 Graze limit and profile predictions (per graze) 1.50 Papers explaining the use of the predictions 2.50

Asteroidal occultation supplements will be available at extra cost: for South America via Orlando A. Naranjo (Universidad de los Andes; Dept. de Fisica; Merida, Venezuela), for Europe via Roland Boninsegna (Rue de Mariembourg, 33; B-6381 DOURBES; Belgium) or IOTA/ES (see below), for southern Africa via M. D. Overbeek (Box 212; Edenvale 1610; Republic of South Africa), for Australia and New Zealand via Graham Blow (P.O. Box 2241; Wellington, New Zealand), and for Japan via Toshio Hirose (1-13 Shimomaruko 1-chome; Ota-ku, Tokyo 146, Japan). Supplements for all other areas will be available from Jim Stamm (11781 N. Joi Drive; Tucson, AZ 85737; U.S.A.) for \$2.50.

Observers from Europe and the British isles should join IOTA/ES, sending DM 40.-- to the account IOTA/ES; Bartold-Knaust Strasse 8; D-30459 Hannover, Germany; Postgiro Hannover 555 829 - 303; bank-code-number (Bankleitzahl) 250 100 30.

IOTA NEWS

David W. Dunham

Total Lunar Eclipse: The most important feature of this issue is the article on occultations during the November 29th total lunar eclipse, the best one for occultations throughout North America in many years; see the following pages.

<u>IOTA/ES Business Meeting:</u> The European Section of IOTA will hold their annual business meeting in Hannover, Germany, on Saturday, November 20, starting at 10:30 am. Contact Hans-Joachim Bode (see the back page) for more information.

ESOP XII: The 12th European Symposium on Occultation Projects was held in Roden, the Netherlands, on August 27-31. Claudio Costa's PC Evans program, along with numerous auxiliary programs and datasets for calculating and distributing detailed (formerly U. S. Naval Observatory, or USNO) lunar total occultations, was distributed to several national coordinators for the first time. At the meeting, I was able to just discuss what needed to be done with the coordinators who were there. Afterwards, I wrote the procedures and updated other documentation to create a consistent and understandable package, a job that took much more effort than I expected and which delayed this issue of **ON** for over a month. See the article on lunar occultation prediction news on p. 17 for more about the "PC Evans package". Many very interesting presentations were given during the main part of the ESOP meeting over the weekend. More about this enjoyable symposium will be published in a future issue. ESOP XIII will be held in late August, 1994, in Cracow, Poland, while ESOP XIV will be held in the Czech Republic in Plzen in 1995, the city's 700th anniversary.

IOTA Annual Meetings: The 11th annual meeting of IOTA was held Saturday, September 11, near Houston, TX; see Rocky Harper's article about it on p. 11. We

tentatively plan to hold the 1994 IOTA meeting in western Texas just before the annular solar eclipse that will occur on Tuesday, May 10. The date would be either Saturday, May 7 or Sunday, May 8, with the location probably either Midland-Odessa or El Paso. Monday, May 9th would be devoted to testing equipment for, and deploying to, the annular eclipse. By selecting a western Texas location, those who also want to attend the Texas Star Party at Prude Ranch near Ft. Davis (and the other meetings that will be held there) could do so, but I think that we want to hold the IOTA meeting at a more accessible location for the benefit of those who are more interested in the eclipse. Let me know your ideas about this (1-301-474-4722 or E-Mail david dunham@jhuapl.edu or see the roster for an address) soon since we want to make a decision by mid-December and include it in the next issue.

IOTA Manual: We plan to produce 100 copies of Wayne Warren's current "Script" version of the IOTA Manual before the end of this year for new observers and for current members who request a copy from the McManuses, to replace the difficult-to-read POM that has been distributed during the last several years. This version still needs some work that I have not had time to do. The "final" version that will be distributed to all members will also need to include revisions to describe the new graze predictions (a small change, since they are so similar to the current ones) and considerably more work to incorporate the "USNO" occultation papers that we have inherited and which are largely out of date.

E-mail: The October 9th occultation by 27 Euterpe was quite successfully observed in the southwestern USA, based on a good astrometric update obtained at Lowell Observatory four days before the event. Electronic mail (specifically, Internet) was valuable for distributing updated predictions quickly to dozens of observers. I mailed (regular postal service) the same information to a couple of dozen other ON subscribers in or near the path for whom I did not have E-mail addresses, but I think at best 25% of those may have arrived in time. In the USA and many other countries, Internet is available at most universities, colleges, government agencies, and large businesses. If you don't have access at work, Internet can be reached for reasonable prices via Compuserve, MCIMail, GEnie, and other vendors in the USA and in other countries; a few IOTA members have already purchased access. Local and regional coordinators are encouraged to gain Internet access to facilitate communication about lastminute updates. Those with E-mail also have a duty to try to inform other IOTA members without E-mail in their areas of local time-critical events.

This Issue and a Correction: This issue marks the first issue of volume 6. With volume 5 issue 12, we exceeded 300 pages by 20 pages, so we decided it was time to start a new volume. We apparently got ahead of ourselves last March, since the cover of issue No. 11 said Vol. 6 when it should have been Vol. 5. The first page of the issue (p. 275) correctly stated "Volume 5".

1994 North American Asteroidal Occultation Supplement: The 1994 N.A.A.O.S., giving Edwin Goffin's predictions, world maps, and finder charts, is being distributed with this issue, and others should have already received similar predictions for their areas from their national or regional coordinator.

The Next Issue: The main purpose of the next issue will be to describe IOTA's planetary occultation predictions for 1994. It will also include several articles, such as Tony Murray's article on new double stars, that could not be included in this issue due to lack of time. We also hope to include Richard Wilds' paper on Watts' lunar limb correction charts, a separate IOTA publication, with that mailing. We should receive your contribution for ON 6 (2) by December 13. We want to produce the issue before the end of 1993, but it might not actually appear until the first or second week of January 1994.

OCCULTATIONS DURING THE TOTAL LUNAR ECLIPSE OF 1993 NOVEMBER 29

David W. Dunham

This eclipse is the best-placed for North America since 1982; the Moon will be in the zenith at mideclipse near Mexico City. But unlike other total lunar eclipses well-seen from the continent during the previous 30 years, several relatively bright stars in the northern part of the Hyades will be occulted. Occultations might be seen during this eclipse from northern and western Europe, northwestern Africa, South America, Japan, eastern Siberia, northern New Zealand, and most of the Arctic, Atlantic, and Pacific Oceans. This eclipse will probably be only slightly darker than usual due to the lingering effects of material still in the atmosphere from the eruptions of Mt. Pinatubo in 1991. Nevertheless, the November eclipse should be very good for occultations due to the brightness of the stars that will be occulted. With large telescopes, occultations of stars as faint as 13th magnitude might be observed in the dark center of the umbra that will be traversed by the northern parts of the Moon. Observers of this eclipse should certainly include the timing of occultations in their program.

Only during a total lunar eclipse can Value: occultation disappearances and reappearances be observed equally well around the entire circumference of the Moon's disk. This gives a rare opportunity to accurately link the eastern and western hemispheres of the Moon. Dr. Chester Watts found this job to be extremely difficult when he constructed his epic charts of the marginal zone of the Moon from photographic plates where in general only either the eastern or western hemisphere were sunlit. Improving information about the lunar profile from lunar eclipse occultation timings is important for all lunar occultation analyses. But this especially benefits analysis of solar eclipse Bailey's head timings made near the edges of the path of annularity or of totality that are used for measuring small variations in the solar diameter. For the solar eclipse analyses, where the most accurate data involve the lunar polar regions, lunar eclipse grazing and near-grazing occultation timings have special value. Since the Moon is near the ecliptic during both lunar and solar eclipses, the latitude libration is always near zero in both cases.

Observing Strategy: The eclipse occurs late Sunday evening-early Monday morning at the end of the Thanksgiving Day weekend for Americans. So if you are travelling during this holiday period, plan to return in time to get ready to observe the eclipse (and possibly to travel further if clouds threaten your main observing location). Try to gain access to the largest telescope possible and concentrate on timing occultations around the entire Moon's limb. Also, try to observe reappearances, as they are often under-observed during eclipses.

If you have a portable telescope, allow youself enough time to travel to view the eclipse. Most parts of North America and Europe are quite cloudy in late November. Mobility could mean the difference between seeing and not seeing this rare event. Remember that Sunday evening traffic will be heavier than usual so allow more time for travelling. If the largest telescope available to you is portable, it would be best to take it to the nearest grazing occultation path. Observations of southern-limit grazes will be more difficult than northern limit since the southern part of the Moon is in the umbra for a shorter time than the northern part. If a southern-limit graze path of a relatively bright star (necessary since the southern part of the Moon will be in the brighter outer portions of the umbra throughout totality) is near you, its observation should have top priority.

Other Publications: Basic information (such as contact times) about this eclipse are given, or can be inferred, from this article. Additional general information about the November eclipse is given on pages 66-

68 of the November issue of Sky and Telescope (S&T). The 1993 occultation highlights article on pages 74-76 of the January issue of S&T gives a map showing 5 eclipse graze paths, double star, and other eclipse occultation information. A reference to this article probably will be given somewhere in the December issue, which will reach (or has reached) virtually all North Americans before the eclipse. I was not able to prepare an article in time for either S&T or Astronomy, because I did not have the software working on my PC to generate the necessary plot for that article until mid-August. This is part of the slow resolution of the problems caused by the loss of my access to two mainframe computers in 1992 September. Astronomy did publish a letter from me on p. 13 of their December issue pointing out the good eclipse occultation opportunity and refering readers to IOTA for more information.

Fortunately, several thousand people who do not get ON will get a chart showing stars to 11th magnitude and other information about eclipse occultations. An article that I prepared will be published in the November issues of the Astronomical Society of the Pacific's Mercury magazine and in the Astronomical League's Reflector. The Mercury article also includes basic information about timing occultations.

The Star Field: The eclipse star field, shown in two charts, is in Taurus about 5° northwest of Aldebaran. The brightest star in the field is 4.4-mag. κ Tauri = ZC 656, near the upper-left corner of the chart. It will be occulted during the final partial phases in the Pacific Ocean southeast of New Zealand. Some points in the northern limit calculated with the OCCULT program are below; since the region of visibility of this occultation is only in the ocean, I presume that nobody will attempt to observe it.

West <u>Long.</u>	<u> Lat</u>	<u>t.</u>	Moon <u>alt.</u>	Sun <u>alt.</u>	umbral <u>dist.</u>	% <u>sunlit</u>
160. 150	-59° -53	32 ′ 44	1 '	-2· -10	90U 94U	66E 72E
$\frac{145}{142}$	-51 -49	00 29	$\frac{14}{16}$		99U 103U	78E 86E

No significant clusters or nebulae are in the star field which does contain several outlying members of the Hyades open cluster.

One plot shows only the stars, down to magnitude 14.0 from the Q-catalog based on the USNO XZ and the Space Telescope Guide Star (GS) Catalogs, the data from the latter which were sent to me via Wayne Warren by Doug Mink at the Center for Astrophysics in Cambridge, Massachusetts. Underlined stars are, or may be, double. The Q-catalog equinox 1950 bounds, which are close to those determined by David Herald,

are 4^h 12.0^m to 4^h 24.0^m in right ascension (R.A.) and +19°8 to +22°3 in declination (Dec.). The equinox 1950 stars-only chart cuts off some stars at the top (which will be traversed by the Moon only as seen at low altitude from the far southern Pacific Ocean where there are no islands), where the declination is $\pm 22^{\circ}2$, and there is a blank strip at the left because the high R.A. needed to end at 4^h 25.0^m. David Herald's equinox J2000 boundaries are 4^h 15.4^m to 4^h 27.3^m in R.A. and -21° 04' to -23° 34' in Dec. "topocentric paths" chart on the page after the "stars only" chart, with apparent-place coordinates so that it can be used with the "USNO" (now IOTA) detailed Ocatalog total occultation predictions, includes the stars that were left out at the top of the 1950 chart, plus a blank strip, but it cuts out a few stars at the bottom (that will be traversed by the Moon only as seen from the Arctic) and at the right (which is shown on the 1950 chart), and leaves a blank strip nearly 1 minute wide at the left (where the Moon will be out of the umbra as seen from anywhere on the Earth's surface).

Topocentric Paths: The other chart shows the same star field, but also includes numbers of the brighter stars, a figure of the Moon generated by Bob Bolster with John Westfall's Moonview program, and topocentric tracks for the Moon's center for 25 locations while the Moon's center is on the chart and is above the local horizon. Apparent place coordinates (with precession, nutation, and aberration applied to 1993 November 29) are used. The Moon's figure is drawn for the right size and orientation during the eclipse. The position angle of the Moon's North Pole (0° of Watts Angle, or WA on IOTA predictions) will be 349°, to help locate reappearing stars with lunar features. In many cases, the pattern of the star field will give a better idea of the point of emersion.

A copy of the Moon figure can be moved with its center along the path, keeping its orientation the same as shown on the chart, to estimate the times and locations of disappearing and reappearing stars. The name of the location, for which a topocentric curve of the Moon's center is plotted, is given along the path. Labels are written near the right (lower R.A.) end of the tracks; if they are not legible, I have also typed them near an uncluttered part of the curve. A list of the coordinates used for calculation of the paths is given in Table 1.

Time increases from right to left (the Moon's R.A. is always increasing) along the curves. If the Moon is above the horizon both times, the paths start at first umbral contact and end at last umbral contact. Tick marks at 1-hour intervals, from 5h U.T. to 8h U.T., extend north from the path. Shorter tick marks extending south of the paths mark the four eclipse

contacts, which will occur at the following times:

h m
4 40.4 First umbral contact (First Contact)
6 02.2 Start of totality (Second Contact)
6 50.1 End of totality (Third Contact)
8 11.9 Last umbral contact (Fourth Contact)

Not shown is mid-eclipse, which will occur at 6^b 26.1^m UT. For stations where the early part of the eclipse occurs before moonrise, the paths end at the following times: Auckland and Khabarovsk, at fourth contact: and Tokyo, at 8h U.T. For paths near the right side of the chart, often the paths start to the right of the chart and end near moonset. These paths can be extended to the right off the chart, but these parts of the paths are shortly after first contact when only the eastern part of the Moon's limb, which is generally then on the chart, is in the umbra. The start and end times for these tracks are as follows: Recife, 5:00 UT to 7:45 UT; La Palma, 5:15 to 7:45; San Fernando, 5:15 to 7:15; Rome, 5:15 to mid-eclipse; Herstmonceux, 5:00 to 7:30; and Moscow, 1st contact to 5:30 UT. For locations not shown on the chart, interpolate.

Table 1. Stations

Location	Latitude	E.Long.
AUCKLAND, New Zealand	-36°908	174°777
TOKYO, Japan	35.660	139.770
KHABAROVSK, Siberia	48.450	135.100
MOSCOW, Russia	55.755	37.570
ROME, Italy	41.924	12.453
HERSTMONCEUX, U.K.	50.871	0.338
SAN FERNANDO, Spain	36.462	-6.200
LA PALMA, Canary Is.	28.758	-17.880
RECIFE, Brazil	-8.051	-34.958
RIO DE JANEIRO, Brazil	-22.898	-43.186
ST. JOHN'S, Newfoundld.	47.537	-52.753
BUENOS AIRES, Argentina	-34.605	-58.434
CARACAS, Venezuela	10.507	-66.928
SANTIAGO, Chile	-33.418	-70.630
MONTREAL, Quebec BOGOTA, Colombia	45.500	-73.600
BOGOTA, Colombia	4.599	-74.081
LIMA, Peru	-12.100	-77.050
MIAMI, Florida	25.750	-80.250
KANSAS CITY, Missouri	38.964	-94.497
MEXICO CITY, Mexico	19.250	-99.100
LOS ANGELES, California	34.113	-118.302
VANCOUVER, Brit. Col.	49.500	-123.100
ANCHORAGE, Alaska	61.210	-149.870
HONOLULU, Hawaii	21.300	-157.850
NORTH POLE, Arctic Oc.	90.000	

The Stars and their Numbers: Apparent positions were plotted so that the RA and Dec. given in the detailed IOTA predictions could be used to locate stars whose occultations are listed. The RA bound was increased by 3^m due to the increase in RA since 1950.

The numbers of stars of magnitude (mag.) 11.0 and brighter have been plotted on the apparent-place chart. The number of each star starts about 3.5 mm to the right of the center of the star; keep this distance in mind when examining the chart, since fainter stars often appear between the star and its number. All star numbers increase with R.A. from right to left across the chart. If available, a three-digit ZC number, ranging from 628 (4.8-mag. ω Tauri, the 2nd-brightest star in the lower right part of the chart) to 657, is plotted. The next preference, if available, is a USNO XZ (or X) number in the 5000's. If neither ZC nor X numbers are available, the star's Q-catalog number, ranging from 93 to 1415 on the chart, are given. Around R.A. 4h 20m, the Q numbers overlap the ZC number range; for example, Q 633 is about 2 cm north of the typed "SANTIAGO" while ZC 633 = 53 Tauri is just above the words "MEXICO CITY". The Bayer Greek letters of the two stars mentioned above, and the Flamsteed numbers of four stars (51, 53, 56, and 67 Tauri) have in addition been typed with larger, darker characters near these bright stars.

<u>Variable Star:</u> The only XZ variable star on the chart is ZC 634 = 56 Tauri = V724 Tauri = Q 536 = SAO 76551, a rotating variable that has an insignificant mag. range for occultation work. The mag. of this "ACV variable" varies from 5.36 to 5.40 in a 1.569-day period. The star will be occulted for observers in southern South America. I have not checked to see if any fainter non-XZ known variables are on the charts.

Double Stars: Known double stars are underlined on the charts. Data about most of them are given in the lists of European and North American grazes given later in this article, while the rest of the visual doubles are listed below. Spectroscopic binaries possibly resolvable by high-speed photoelectric occultation recordings, that are not in the graze lists, include X 5728 (SAO 76609) and ZC 665 (SAO 76618).

Visual Doubles not in Graze Tables


<u>0500#</u>	Ľ	JAC/BD	Desig.	Иазі	Maul	<u>Jep.</u>	<u>P.A.</u>	3
X 15 6 3 3 X 15 6 3 3	Ā		ADS3118 SI (4328	1 4.2 1 4.0	7.5	1.5	[1] F F	
$(f_i)^* = f_i \stackrel{E_i}{\leftarrow} \mathcal{E}_i$	$ \forall $		HIT (44.1%)	50.0	i,	1.	115	
C. 12 11 18			A COURT OF A COURT				2.1	

Mag1 is the magnitude of the primary, Mag2 is that of the secondary, Sep. is the separation in arc seconds, and P.A. is the position angle. The star's double-star code is given after the USNO#, under D. Under SAO/BD, 5-digit numbers are SAO numbers; one of the stars is a BD star that is not in the SAO. Under Desig. (designation), ADS is Aitken's Double Star, CL is C. Lynns, who discovered the star with speckle interferometric observations at Kitt Peak Nat. Obs. in 1975 October, SI is B. Sincheskul, Poltava, Ukraine, who saw a gradual fade when the star disappeared on 1974 March 28, and HP is Harold Povenmire, Florida, a confirming observation of occultation duplicity obtained on 1974 April 25 (the star is K Tauri). Note 1: X05583's northern limit crossing the Bahamas is shown as track #1 on the North American graze map, but since the secondary is fainter than mag. 12.5, duplicity information was not included in the XZ catalog, so it has no double star code (blank) in the IOTA-USNO total occultation and graze predictions. Note 2: I did not know about Q 1204's duplicity, so its code is also blank in the Q-catalog prediction input file for calculating detailed total occultation predictions with the Evans program. Like nearby K Tauri, the star will be occulted in the umbra only for part of the far southern Pacific Ocean.

Minor Planets: Using data in Ephemerides of Minor Planets for 1993, I found only one asteroid that will appear on the charts at the time of the eclipse. It is 15.5-mag. (2003) Harding, which is labelled on the topographic paths chart to the right of "AUCKLAND". Its occultation will be visible in the umbra only from part of the Pacific Ocean east of New Zealand.

November Version of the USNO Q-catalog: The first edition of the Q-catalog was described in ON 5, 208, and updates made for last June 4th's eclipse were described on pp. 302-303 of the last issue. The range of Q-numbers for the November 1993 lunar eclipse field is 5 to 1427.

I recently manually compared a B1950-version of the appropriate part of the Bonner Durchmusterung (BD) with the Q-catalog to determine the BD numbers for many of the non-XZ Q-catalog stars. These BD numbers are not in the predictions because the database (Besselian elements file, or befile) used by the prediction program had to be created over a year ago at USNO before thier IBM mainframe computer was taken away. These BD numbers are listed in Table 2.






Table 2. BD Stars in the Q Catalog

Q No.	B.D. No.	GSC No.	Q No.	<u>B.D. No.</u>	GSC No.
Q 5 Q 55 Q 109 Q 141 Q 228 Q 228 Q 228 Q 230 Q 306 Q 323 Q 389 Q 415 Q 434 Q 481 Q 633	+20° 718 +20 719 +19 683 +20 722 +20 723 +19 686 +21 613 +19 688 +20 726 +21 615 +20 728 +19 693 +20 730 +21 620 +20 732 +21 620 +21 624	12630981 12631027 12590195 12720234 12760341 12760254 12760254 12760375 12760940 12760399 12720180 12761221 12761074 12760222 12761271 12760280 12760280 12760280	9 558 9 673 9 688 9 715 9 716 9 745 9 755 9 819 9 881 9 1050 9 1063 9 1080 9 1117 9 1134	+19 * 702 +20 737 +19 703 +21 627 +20 738 +20 739 +21 628 +21 629 +20 742 +20 743 +20 743 +20 747 +20 752 +21 638 +19 711 +19 712 +19 714 +19 715	12720345 12760325 12720683 12761368 12760246 12720572 12761438 12760840 12720728 12761556 12720945 12761505 12761505 12761284 12720767 12730688 12730379
Q 655	+22 677	12761439	Q1244 Q1415	+19 719 +21 646	12730747 12770282

The star $+21^{\circ}$ 634 was noted as "doubtful" in the BD, and no star brighter than mag. 12.7 was found near its position, so no Q-catalog star has been given that BD number. The double star code of X 5678 = Q 1054 was incorrectly given as "D" in the earlier versions of the Q-catalog. X 5678's correct BD number is $+21^{\circ}$ 638a; BD $+21^{\circ}$ 638 = 9.4-mag. Q 1050, which is more than 20' north of X 5678.

I found five duplicates in this part of the Q-catalog, all apparently caused by problems with matching when I computer-merged the XZ with the Q-catalog over a year ago. In each case, the XZ data should supercede the unmatched Q-catalog data, as shown in the table below. Unfortunately, the five now-rejected stars were included in the Q-catalog befile mentioned above, and therefore continue to be included in the Q-catalog detailed occultation predictions.

Action on Q-Catalog Duplicate Stars

Rejec Star	ted mag.	Retained Q No. ZC/XZ mag.
Q 88	9.5	Q 87 X 5519 10.0
Q 106	7.7	Q 105 X 5522 8.1
Q 198	8.4	Q 197 X 5534 8.9
Q 753	6.5	Q 751 ZC 642 6.9
Q1004	8.3	Q1003 X 5666 9.6

<u>Predictions:</u> The Q-catalog predictions and their distribution were described in **ON 5** (8), p. 20 and (9), p. 240. More detailed information is in the verification form explanation that was distributed with those predictions. For those without predictions, use the star field charts here to locate both disappearing and reappearing stars, and try to mark the ones without an identifying number. Accurate Q-catalog predictions can be computed to identify the stars after the observations are made. If time permits, they might also be generated before the eclipse upon request to a regional

coordinator who has the PC-Evans program; see the "Lunar Occultation Prediction and Software News" article later in this issue.

Reporting Observations: Occultation timings during this eclipse should be reported on the International Lunar Occultation Centre (ILOC) lunar occultation report forms, or the equivalent IOTA/ILOC graze report forms, or in an ASCII file on MS-DOS-compatible diskette [for the latter, see ON 4 (10), p. 237 and ON 4 (5), pp 92-97]. For all occultations that occur during lunar eclipses, please also send a copy of your report to David Herald; P.O. Box 254; Woden, ACT 2606; Australia. He will analyze all timings made during the eclipse and publish his results in ON. For the star number, use the ZC number and catalog code (column 16) "R". If the star is not in the ZC, give its SAO number and put "S" in col. 16. If it is in neither the ZC nor the SAO, give its X number with "X" in col. 16. If the star is in neither of these catalogs, give the star's Q-catalog number, if you have Q-catalog predictions, and put "Q" in col. 16. If you don't know any of these numbers for an observed occultation. include a copy of the star chart with your report marking these fainter stars whose occultations you time.

Grazing Occultations: Eberhard Riedel, Munich, Germany, prepared maps and tables of all grazes of XZ stars that will occur against the umbra during this eclipse. He prepared these maps in the same style as those in the hemispheric grazing occultation supplements for 1993 distributed earlier. But rather than prepare maps covering large parts of each hemisphere. like those published on pages 304 and 305 of the last issue, Riedel produced more legible maps only covering the more populated parts of North America and Europe that are included, along with their associated tables, on the following pages. Another map (and associated table) was prepared for South America and is being send only to ON subscribers there. Paths end in a "B" when they occur at the edge of the umbral shadow. Points along the paths where totality begins and ends are not indicated, but you can tell whether the eclipse occurs during totality or during a partial phase from the time of the event in your area. On the map of Europe, the altitude limits are too lenient.


<u>Lunar Polar Diameter:</u> Two occultations offer reasonable possibilities for observation at both the northern and southern limits for an accurate lunar polar diameter measurement. Preliminary information was given about them on p. 306 of the last issue. That information is reviewed and updated here.

<u>ω Tauri, N. Limit:</u> At mag. 4.8, ω is the 2nd-brightest (not the brightest, as stated last time) star that will be occulted during the eclipse, around 5:30 UT,

during the early partial phases in Europe. The northern limit passes 5 km n. of Alesund and 70 km n. of Oslo, Norway; 35 km n. of Motala, Sweden (Alt. 11°); 10 km s. of Vilnius, Latvia (Moon alt. 5°); and near Minsk, Belorus (but Moon alt. only 2°). Eberhard Bredner hopes to travel to Scandinavia to help local efforts to observe this graze.

<u>ω Tauri, S. Limit:</u> During totality around 6:15 UT, the southern limit passes 20 km south of Villa Cisneros, Southern Sahara, and north of Atar, Mauretania (Moon alt. 12°). Hans Bode is looking into the possibilities for an IOTA/ES expedition to

these areas, but he does not expect to go there himself; he will probably go instead to the Canary Islands to observe the occultation of a star by Jupiter that happens to occur during this eclipse (however, the Canary Islands are well-placed for a valuable southern-limit graze of 8.7-mag. $\times 5548 = \text{SAO } 76533$, track 9, whose northern limit, track 8, is far north in Scandinavia). Nevertheless, an expedition to the Villa Cisneros or Atar area for the spectacular graze of ω Tauri would be very interesting and valuable, as well as a good escape from the cold and likely cloudy weather in Europe.



North American Grazing Occultations during the Lunar Eclipse of 1993 November 29


<u>No.</u>	Year Mo Day US	NO SAO D Ma	g %Snl L		ong Lat Star Name	Mag1 Mag2 Sep PA
1	1993 Nov 29 X 55	83 76546 8.	2 62E N	n m 58.7	92° 23°	
2	1993 Nov 29 X 55	90 76550 8.	7 62E N		104 36	
3	1993 Nov 29 6	33 76548V 5.	4 61E N	5 9.1 1	101 32 53 Tauri	Spectroscopic
4	1993 Nov 29 6	42 76564K 6.	9 59E N		140 32 219 B. Tauri	7.7 7.7 0½10 353°
5	1993 Nov 29 X 56	24 76565 7.	1 58E N		40 21	0,10 333
6	1993 Nov 29 X 56	53 10.	0 30E N		36 15	
7	1993 Nov 29 6	46 76571M 6.	1 30E N	6 16.2 1	140 54 224 B. Tauri	6.2 9.3 1.90 170
8	1993 Nov 29 6	46 76571M 6.	1 10E S		24 21 224 B. Tauri	6.2 9.3 1.90 170
9	1993 Nov 29 x 56	58 76579 9.			40 26	0.2 9.3 1.90 170
10	1993 Nov 29 x 56	60 76580 8.	3 19E N		40 16	
11	1993 Nov 29 X 56	61 76581 8.	5 17E N		40 29	
12	1993 Nov 29 X 56	55 76578 9.	1 6E N		40 58	
13	1993 Nov 29 6	51 76585x 5.9	9 6E N		40 45 227 B. Tauri	6.7 6.7 0.15 55
14	1993 Nov 29 X 56	14 10.1			00 47	0.7 0.7 0.15 55
15	1993 Nov 29 X 56				40 52	

ZC 646, N. Limit: The northern limit for 6.1-mag. ZC 646 passes 65 miles n. of Ketchikan, AK; 20 mi. s. of Ft. Nelson, Brit. Col.; 7 mi. n. of Meander River and over Ft. Fitzgerald (where totality ends at central graze), Alberta; 40 mi. s. of Ft. Chimo, Quebec; and 25 mi. s. of Nain, Labrador.

Members of the Edmonton center of the Royal Astronomical Society of Canada are making extensive preparations to observe the graze. Russ Sampson is leading the effort, with help from IOTA member Doug The observers would travel by bus to Ft. Nelson, and rent winterized vehicles to reach the graze zone, where special precautions will be needed to cope with temperatures that may be below -20°. There is about a 40% chance that the sky will be clear or have scattered clouds, and not too windy to prevent observation. The effort is expected to cost \$2000, 10% of which IOTA has agreed to cover (and half of that a private donation); the organizers are optimistic about getting the remainder covered by a Scientific Research Council grant. An article about these preparations was published in a recent RASC bulletin.

ZC 646, S. Limit: During totality, this passes about 150 km northwest of La Paz, Baja California, north of where Don Stockbauer and I observed a graze of Atlas during the 1991 July 8th Pleiades passage. Southern Baja California is a free-trade zone with no customs inspections, facilitating the temporary inport of telescopes and other equipment. Paul Maley plans to observe this graze with at least one other observer: contact him (1-713-488-6871 or see the roster for an address) if you might be interested in helping this effort. The path crosses Highway 1 on the west side of Baja, where marine cloud may be a problem. There will likely be no clouds where the path crosses the east coast of Baja, but that seems to be accessible only by boat. The expedition for this graze also expects to get some GPS measurements to resolve a large map discrepancy that was found at the Atlas graze sites.

Joe Senne calculated details of these grazes.



November 29	Duplicity				Spectroscopic	· v							Spectroscopic
European Grazing Occultations during the Lunar Eclipse of 1993 November 29	Lat Star Name		17°	45	70 Omega Tauri	29 Omega Tauri	1	30	34	70	33	28	22 53 Tauri
mar Ec	5 0			15	20	30	v	30	9	-16	59	30	30
g the tu	SSnl L W.U.T. L	E	5 8.9	5 1.0	5 9.2	6 2.4	5 17.7	6 7.0	6 8.2	5 14.7	6 0.2	7 9.8	7 10.4
Ĕ	الب		2	*	×	s	×	cs	z	2	S	2	z
s dur	Snl		72E	72E	68E	3ξ	389	ZE	67E	- 889	ZE.	62E	61E
ation	ag ag		ø.		4.8	4.8	9.3	9.3	10.0	8.7	8.7	8.7	5.4
g Occult	SAO D Mag		76524	76525	76532K	76532K	76534	76534		76533	76533	76550	76548V
Grazin	ONSO		x 5518	X 5522	628	628	K 5551	< 5551	6255 x	× 5548	£ 2248	(5590	633
a	≥ı		Ò.	3	0	0	0	<u>o</u> .	0	φ.	0	0	ō.
ğ	Mo Day												
ä	鈻		2	ŝ	£	£	ŝ	ž	ŝ	£	읖	£	ŝ
_	Year		1933	83	1993	1833	1993	1993	1833	1893	1993	1993	1993
	2		-	7	M	4	Ŋ	•	~	∞	٥	10	Ξ

Other Graze Expeditions: Probably the most spectacular graze in the USA will be seen near the northern limit of the occultation of 5th-mag. 53 Tauri, especially in North Carolina, where the graze occurs shortly after totality begins. But the southern limit crosses remote and dangerous parts of Colombia and the Amazon basin. There are similar problems with the other limit for all grazes in the contiguous USA. Bob Stewart, Norfolk, VA, is planning an expedition to the vicinity of Jacksonville, NC, and there will probably also be an expedition for 53 Tauri near St. Pauls, close to highway I-95 south of Fayetteville, NC. Harold Povenmire plans to observe this graze near Water Valley or Courtland, MS, where another northern-limit graze, of 7th-mag. SAO 76565, can also be seen. The RASC Edmonton Centre, as part of their training for the ZC 646 graze, observed a northern-limit graze of 53 Tauri near Edmonton on September 8th. analysis of those observations should improve the prediction for the November 29th occultation of that star, and will probably improve the predictions for other northern-limit grazes as well. The results of this work will be put on the IOTA occultation line answering machine at 1-301-474-4945, along with weather updates and plans for other expeditions; please give me the latter information if you are planning a graze expedition, preferably by telephoning me at 1-301-474-4722.

THE 11th ANNUAL IOTA MEETING

Rocky Harper

The 1993 annual meeting of the International Occultation Timing Association was held on September 11, 1993 at the Lunar and Planetary Institute, Houston, Tx. The meeting began at 9:00am with the call to order by president David Dunham. Nine members were present, including the Dunhams from Greenbelt, MD.; Dan Falla from San Diego, CA; Rick Frankenberger and Gary Dowdle from San Antonio, TX; the others were all from the Houston, TX, area. Visitors from the Houston Astronomical Society and the JSC Astronomical Society also attended.

The first order of business was the financial report which showed IOTA to be healthy. The largest costs are printing and mailing the ON. No increase in membership dues is seen at this time. Work on the IOTA Observer's Manual has continued with a release hopefully in 6 months. Next David mentioned that he thought that IOTA should reincorporate in Maryland or Kansas. IOTA first incorporated in Texas on August

19, 1983 which meant that the organization had by-laws and a charter. It obtained tax-exempt status from the IRS, as long as there was an annual meeting in the state of incorporation. After 10 years in Texas, David feels it's time to consider moving the meetings to another area.

Next was a discussion of total lunar occultation prediction software. David talked about the PC EVANS program and David Herald's OCCULT program. See the May 1993 ON for more details. A crude version of the Evans program (it just applied linear a and b factors to standard station data that were keypunched) was originally written for a mainframe computer by Carroll Evans of China Lake, California back in 1962. The current sophisticated version was written by Tom Van Flandern at USNO in 1970, and updated several times to add new features since then. It was converted to PC use by Claudio Costa in Rome a year ago. David brought his PC which had the PC EVANS and OCCULT programs on hard disk which some members copied. The improved coverage of lunar graze predictions and maps by Eberhard Riedel was also discussed. The mention of the observation of unpredicted events brought on a good talk about the future improvement of predictions. Star position errors should be virtually eliminated as well as better magnitudes derived when data from the Hipparcos space mission are reduced. David also showed how the "25 error in the Watts data should be reduced to "1" using information from the upcoming Clementine Moon/asteroid mission scheduled for launch in early 1994. David is personally involved in this mission so it should be exciting. David next gave a very interesting talk about the 12th European Symposium on Occultation Projects that was held in Roden, the Netherlands, on August 27-31. Then we broke for lunch to a local cafeteria where we continued our discussions.

The meeting resumed around 1:00pm with the highlight on video developments. A lot of talk focused on Tom Cambell's VTACT system and the European Cuno video time inserter. Image intensifiers were also discussed. Gary Dowdle mentioned that relatively inexpensive video couplers for image intensifiers are advertised in law enforcement, but not astronomical, publications. He was going to do some research on this and write something about it for ON.

The next item concerned lunar and solar eclipses. Much was said about the November 29, 1993 total lunar eclipse at which time it passes through the Hyades star cluster producing some excellent occultations. Future expeditions were mentioned including those to the May 10, 1994 annular solar eclipse here in the United States. The next total solar

eclipse occurs November 3, 1994 in South America where several expeditions are planned.

At 3:15pm David talked about upcoming asteroidal occultations and major grazing events. The last part of the meeting was used to plan for a southern-limit graze of 4.3-magnitude Alpha Cancri (Acubens = ZC 1341) by a 10% sunlit waning Moon the morning of September 13th. Unfortunately, it was too cloudy to observe that event and the expedition was cancelled.

As always the meeting was very interesting and exciting. There was the usual enthusiasm and a lot of great stories told.

[Ed. We thank Rocky for this report, written in spite of a personal disaster that occurred the ill-fated morning of Sept. 13: A gas explosion destroyed half of his house and burned his arms and face! Fortunately, Rocky had insurance, made a good recovery, and none of the other members of the household were hurt. He wrote this report as soon as his PC was replaced.

1994 LUNAR OCCULTATIONS OF PLANETS

The maps showing the regions of visibility of lunar occultations of major and bright minor planets are reprinted by permission from the Japanese Ephemeris for 1994, published by the Hydrographic Department (which also includes the International Occultation Centre, ILOC) of the Maritime Safety Agency of Japan. On the maps, in region 1, only the reappearance will be visible; in region 2, the entire occultation will be visible; and in region 3, only the disappearance may be seen. Reappearances occur at sunset along a dashed curve, while disappearance is at sunrise along a curve of alternating dots and dashes. We have added the lunar percent sunlit to each map. Some of the maps are given below, while others are in convenient locations on later pages of this issue. Consult these maps especially for occultations of minor planets, which are not included in the detailed IOTA total occultation predictions (occultations of major planets are included).

Those interested in observing partial occultations, which occur within a few kilometers of the northern and southern limits of these occultations, should request predictions preferably three months in advance from Joseph Senne; P.O. Box 643; Rolla, MO 65401; USA; telephone 1-314-3646233; Internet c0458@umrvmb.umr.edu.


TREASURERS' REPORT

Craig and Terri McManus

Our report for the period September 1, 1992 to August 31, 1993 is as follows:

INCOME	
Full Memberships	\$5,765.00
Gifts from Members	13.50
Interest on Checking	93.74
Non-member Predictions	7.50
ON Subscription Only	1,020.00
ON Back Issues	25.00
Prelim Occultation Manual	7.50
Sale of IOTA Items	36.01
Other	3.84
4	
TOTAL INCOME	\$6,972.09
EXPENSES	
Ast, and Graze Supplements	\$860.06
Credit Card Costs	33.46
Mailing Costs (all)	3,489.09
ON Costs	2,452.25
Office Expenses	168.04
Other Expenses	343.38
USNO Total Occ Expenses	234.07
Other	203.24
TOTAL EXPENSES	\$7,784.49
OVERALL TOTAL	-\$812.40

Our current balance is \$3,447.91. The expenses and incomes look a little skewed to the negative due to the fact that we produced five (5) ON's this last fiscal year as opposed to the normal four.



1994 SOLAR ECLIPSE PUBLICATIONS

Since 1949, USNO published a special series of free circulars containing detailed information about important upcoming solar eclipses. Funding cutbacks forced USNO to discontinue these circulars in 1991; the last of these circulars described the 1992 January and June eclipses.

Fortunately, a similar series has been started by Fred Espenak and Jay Anderson. NASA Reference Publication 1301 (April 1993) gives information about the 1994 May 10th annular eclipse and NASA Reference Publication 1318 (October 1993) does this for the 1994 November 3rd eclipse. The format of these publications is similar to that of the USNO eclipse circulars, but some additional information is included, including (of special interest to ON readers) tables of corrections to the northern and southern limits to define "graze zones" based on Watts' profile data. A drawback is that the table of mapping coordinates for the umbral path is given at only 1°-intervals of longitude, making quadratic interpolation necessary for accurate work, and the maps are at 1:5,000,000-scale, 2.5 times less detailed than those in the recent USNO circulars. On p. 16 of Publ. 16, an old address is given for the USGS (the Arlington, VA office was closed a few years ago) and no address is given for Canada. The correct addresses are:

Branch of Distribution
U. S. Geological Survey
Box 25286, Bldg. 810 Federal Center
Denver, CO 80225
USA (or telephone 1-800-USA-MAPS)

and

Canada Map Office 615 Booth Street Ottawa, Ontario, K1A 0E9 Canada

On that same page, the old address for IOTA in Topeka is given (see the top of p. 1 of this issue for the current address) and it failed to give Dr. Fiala's address at USNO for receiving observations (that oversight was corrected on p. 18 of Publ. 1318).

Robert Victor at Abrams Planetarium, E. Lansing, MI, points out a discrepancy in Table 6 of Publ. 1301: The interior and exterior corrections, for both the N. and S. Limits, seem to have been reversed. Robert will communicate with Fred Espenak to try to resolve this discrepancy, and we will mention the result in the next ON. The similar data in Table 6 of

Publ. 1318 seem to be correct.

Those planning expeditions to the limits of annularity or totality are encouraged to obtain these publications; it will probably be a few months before Alan Fiala at USNO and I can calculate more detailed limit data for eclipse planners. Free copies of these publications can be obtained from

Jay Anderson Prairie Weather Centre 900-266 Graham Avenue Winnipeg, Manitoba, R3C 3V4 Canada

Internet: janders@ccm.umanitoba.ca

Requests should include the following information: Name, size, type, and activities of organization; name of contact person; complete postal address; and whether publications are wanted for a) specific eclipse (give date), b) eclipses with local interest only, or c) all eclipses.

Additional useful information about the 1994 November 3rd eclipse is given in good articles in this month's issues of S&T and Astronomy, and more information on the 1994 May 10th event is expected in the 1994 January issue of S&T.

SOLAR ECLIPSE NEWS

David W. Dunham

Annular Eclipse, 1994 May 10: See the previous article and the "IOTA Annual Meetings" subsection of IOTA NEWS on p. 1.

Total Eclipse, 1994 November 3: Paul Maley informs me that his expedition to Tacna, Peru, described on p. 294 of the last issue, is sold out. Unfortunately, everyone in his expedition wants to observe near the center line near Tacna, with nobody willing to stay behind at Arequipa to observe a short distance inside the northern limit of totality. However, if anyone wants to try that, they should contact Paul (1-713-488-6871), who felt that suitable arrangements to observe near the northern limit in the Arequipa area could probably be arranged. However, travel to and from Arequipa would need to be arranged independently of his expedition.

Tom Van Flandern has issued "Eclipse Edge Expedition '94 Bulletin #1 -- July, 1993" giving preliminary information about his planned expedition to a location about 5 km north of the southern limit in the Atacama

Desert near Arica, Chile. The expedition will last from Oct. 28 to Nov. 4 and will include astronomical lectures, night observing sessions of the southern sky, and recreational side trips. The cost from Miami will be \$1700, or somewhat less per person for 2 or 3 sharing a hotel room. A final commitment date of 1994 July 15th has been set. The bulletin, besides giving further details about these arrangements, also has a good explanation of the advantages of locations just inside the path edges over locations near the central line. To get a copy of the Bulletin write to Eclipse Edge Expeditions; PO Box 15186; Chevy Chase, MD 20825-5186. Eclipse Edge Expeditions is also selling, for \$20, a video tape of the successful 1991 July 11th 1st Eclipse Edge expedition in Mexico. including some of my footage of that event.

Hans-Joachim Bode is organizing an IOTA/ES expedition to Bolivia to videorecord and time Bailey's bead phenomena near both limits of this eclipse.

VTACT

Tom Campbell, Jr.

I am now building copies of the VCR Time signal Audio conditioner and Clock Trigger (VTACT), discussed in **ON 5** (8), p. 216. It does the following useful jobs:

- 1. Eliminates microphone and preamp mixer combination.
- 2. Includes a speach filter to enhance voice recording.
- 3. Includes a 1000-Hertz and 1200-Hertz tone filter to greatly enhance the WWV or WWVH minute tone, especially valuable for reliably triggering a video time inserter; as long as you can hear the tone at all, it will trigger properly.

The unit is very stable, with a delay jitter of only ± 2 msec. The unit weighs 1.5 lbs. or 680 grams and fits in a 6 x 5 x 3-inch box. I have one unit, which has 3 printed circuit boards, for sale for \$250. Future units will cost somewhat less since they will use an easier-to-construct single-board design. For more information, contact me at 5405–98th Ave.; Temple Terrace, FL 33617; phone 1-813-985-1842.

GRAZING OCCULTATION OBSERVATIONS

Richard P. Wilds

This quarter's article starts out with a little personal information. I have been transferred. I will still be teaching in the same complex as Craig McManus, but I will have a different office with a different work phone - (913)296-4343. One advantage is that callers can now leave messages with the receptionist, if they cannot contact me directly.

The graze report starts with two reports sent in by Henk Bulder from the Netherlands. The 08/24/92 graze was discussed in detail in the March 1993 ON 5, #11, pages 277-280. The 12/05/92 graze was in the Cassini Region among the polar mountains M4 and M5. Since you will have received your copy of The Marginal Zone of the Moon. I hope the reader will have little difficulty in locating the regions in which the reported grazes were observed [Ed. Unfortunately, distribution of this will be delayed to the next ON].

The first graze of 1993 reported for this quarter was the only graze from H.A.R.T. This graze was also in the Cassini Region. Another early graze was reported from Germany. This was a difficult marginal graze which was not included in Mr. Buttner's regular graze predictions. We also received a report of a well observed graze from Pierre Vingerhoets, president of the Occultation Group of the Belgian Association for Astronomy. The graze was observed in an area south of the crater Sylvester.

The same weather conditions which have given H.A.R.T. difficult, and even dangerous, flooding conditions in the plains have helped observers in the east. Mike Kazmierczak reported a graze observed from Georgia on 02/02/93 - the first of four in the list. All his grazes were very successful, but his best was on 07/10/93. He set up next to a railroad crossing. He reports observing 18 events! This alone would get the heart going, but a train came moments after the last event! That would be a tape to save.

Two midwestern observers also had fine grazes to report. Robert J. Modic had a miss around the crater Brianchon, but he finished with two very good grazes around the crater de Sitter. Wayne Hutchinson also had a graze around Brianchon. These two gentlemen were joined by Robert H. Hays Jr. who reported two more grazes in the Brianchon area. Mr. Hays reported that his 5/27/93 observation of Alpha Cancri was quite an eyeful. He reports several extremely interesting events of a gradual nature which gave the impression of viewing the graze in "slow motion." Other observers have reported such observations and all are left with a feeling of awe at what they have seen.

John Centala must be commended for leading a graze in the middle of "Lake Iowa." He reports that the graze would have been even better if clouds had not been involved.

Perhaps the best report of the quarter also came from a most interesting area of the astronomical world. Poland has a long and proud history of greatness in astronomy. The latest representatives of this tradition are Marek Zawilski and Janusz Słusarczyk of the Section of Observations of Positions and Occultations of the Polish Amateur Astronomers Society. Zawilski writes that the team he works with has had many cases of lost grazes due to clouds and site failures for various reasons. They struggle with more problems than the average graze team due to difficulties such as poor map coverage. He reports that the current effort was actually carried out by two teams -one from Lodz and Warszawa led by Mr. Zawilski and the other from Krakow led by Mr. Slusarczyk. For many of the observers this was their first viewing of a graze. They were very satisfied! This is a tribute to years of perseverance and to overcoming great difficulties in the practice of good science. They correctly identified their graze area as falling between the craters Merril and Rozhdestvenskiy.

Two Florida observers make a return this quarter. Tom Campbell reported a video of a graze in the same area observed by his Polish colleagues. Hal Povenmire also reported observations in this area.

Joaquim Garcia made another report of a graze too difficult for his current IOTA predictions. He reports, though, that he is working on obtaining the software he needs to make his work more valuable to the program. He also reports Rui Goncalves has been helping him with the graze effort for years and had an important part in the current results. It is interesting that these two gentlemen from Portugal joined Mr. Zawilski from Poland to assist Mr. Costa and Mr. Bulder in the first graze on our current list.

HISTORICAL FLASHBACK. A new feature to appear from time to time is a look back at some of historical moments in the observations of occultations. I would be interested in any thoughts and ideas for this section.

For this flashback we return to Westchester, Connecticut on 30 May 1966. Graze leader David Dunham arrived at a secluded site to observe the graze of 6.8 mag. SAO 139072 (ZC 1864). He reports: "The owners of the house figured that we could have found them only with the aid of photographic satellites! While setting up my telescope near a farm house, a small boy from the house said, 'Can I observe it, too?' I turned around to see that he had set up a telescope of his own!" Dave, you can certainly build graze teams in the strangest ways.

Graze List

UTDate V			%					# # S Ap		Аp		N		
YYMMDD	P	Star #	Mag		CA	Location	Sta	Τm	\$	Cm	Organizer	Sh S	WA	8
1992	-								_					
920824	v	78297	2.8	22-	1.0N	Minturno, Italy	201	166	1	5	C.Costa/H.J.J.Bulder	0.4N	0~	1:3
921205	•	92395				Wieringerwerf, Neth.					Henk J.J. Bulder	0.0	177	-5.3
1993														
930130		92600	9.0	41+	3.38	Burlington, Kansas					H.A.R.T. C.McManus	0.0		-4.9
930201		76430	4.5	68+	0.0N	Stollberg, Germany					D.Buttner/A.Viertel	?	_	-1.4
930202		76608	4.3	71+	10.0N	McDonough, Georgia	_				Mike Kazmierczak	0.18		-1.2
930227	٧	93237	8.0	32+	10.0N	Tisselt, Belgium	5				Pierre Vingerhoets	0.15		-2.7
930301		76430	4.5	44+	11.0N	Barnesville, Georgia	1				Mike Kazmierczak	0.48		-1.4
930331		78632	8.1	49+	13.2N	Highland Heights, OH	1	-				>0.5s		3.3
930427		78391	8.4	25+	4.2N	Clodine, Texas					Wayne Hutchinson	0.0		3.2
930427		78394	8.0	25+	10.3N	Schneider, Indiana	1				Robert H. Hays Jr.	0.38		3.3
930427		78395				Cascade, lowa	_		_		John Centala	0.78		
930428		97087	6.8	37+	15.0N	Bundaberg,Qld,Austra	. 1					>0.3 S		
930429		98267	4.3	53+	9.4N	Bolimow, Poland	_			_	Marek Zawilski	0.0	10	
930429		98267				Lubaczow, Poland					Janusz Słusarczyk	0.0	10	6.6
930502	٧	118634				Bradenton, Florida	2				Tom Campbell	0.58		7.3
930510		187498				Richmond Hts., Ohio	1	-	_		Robert J. Modic			-2.7
930514		145965				Richmond Hts., Ohio	1	-			Robert J. Modic			-6.0
930515		146415	6.4	32-	4.0N	Conyers, Georgia	1	_			Mike Kazmierczak			-6.1
930527		98267	4.3	31+	8.2N	Grand Ridge, Illinoi	s 1				Robert H. Hays Jr.			6.7
930527		98267	4.3	31+	7.4N	King, North Carolina	_	_			Hal Povenmire			6.7
930613		128538		42-		Quebradas, Portugal	_				J.Garcia/R.Goncalves			-5.7
930710		128487	6.9	65-	10.0N	Acworth, Georgia	1	18	3 1	25	Mike Kazmierczak	0.2N	349	-5.9
	-								-					

REMEMBER to apply the following shifts:

- 1. Northern limit, waxing-phase, dark-limb grazes tend to have a 0"3 south shift from your predicted graze path. One should spread out, however, since star errors could increase this shift or reduce it to a 0" shift.
- 2. Southern limit, waxing-phase, dark-limb Cassini region grazes tend to have a 0".4 0".5 second of arc south shift from your predicted graze path. Cassini region grazes have profile points from 3 to 7. Southern-limit Cassini grazes will also have large negative latitude librations. This correction should continue into waning-phase grazes to Watts angle 187°.

Please report all grazes to:

Richard P. Wilds 3630 S.W. Belle Ave Topeka, KS 66614-4542 USA

and to the:

International Lunar Occultation Centre (ILOC) Geodesy and Geophysics Division Hydrographic Department Tsukiji-5, Chou-ku Tokyo, 104 Japan

Stellar	cross	reference:
SAO #	ZC #	other
78297	976	Ц Gem
92395	197	,
92600		x2420
76430	599	37 Tau
76608	660	69 Tau
93237	446	
78632		x9576
78391		X9083
78394		X9100
78395	989	15 Gem
97087	1151	
98267	1341	α Cnc
118634	1605	62 Leo
187498	2757	36 Sgr
145965		x30506
146415	3371	
128538		x32200
128487	3524	

LUNAR OCCULTATION PREDICTION NEWS

David W. Dunham and Walter I. Nissen, Jr.

Detailed Total Occultation Predictions: The International Lunar Occultation Centre (ILOC) in Tokyo has computed most of the total occultation predictions for 1994 using their own program, and they will mail the predictions early this month. For their main prediction run, they used the station and prediction options data updated by us over a year ago for the 1993 predictions, which we ran with the EVANS program mainly at USNO before their IBM mainframe computer was taken out of service on 1992 October 1. We have finally updated these input datasets for the 1994 predictions, using information supplied by observers on their 1994 prediction verification forms, and sent them by E-mail (via Mitsuru Sôma at the Japanese National Astronomical Observatory, Mitaka, Japan, since ILOC does not have E-mail) late the first week of this month. At their request, we provided ILOC with the 1993 prediction address list in electronic form (again via Sôma) at the end of October; a version including the 1994 updates was sent late the first week of November. We have asked ILOC to provide 1994 predictions for the new observers, and for new and updated stations, that are in our new datasets but that were not in the 1993 prediction datasets that they have had for almost a year.

According to Dr. Yoshio Kubo of the Japanese Hydrographic Office (of which ILOC is a part), ILOC's prediction program "is the same as the one by the EVANS program and the predicted times differ by only one or two seconds at most". ILOC's program probably produces comprehensive predictions of observable occultations of XZ stars, which is almost 99% of what the EVANS program does. But we are not sure that ILOC's program has all of the features of the EVANS program, including observability code rejection based on IOTA's modification of B. Schaefer's MAGLIM subroutine (or observability-code scheme), "graze nearby" messages, photoelectric-option output, occultations of K-catalog stars and galactic-nebular objects, occultations of major planets, terminator-distance messages when appropriate, and events during umbral lunar eclipses (in 1994, there is only a shallow eclipse centered over southern Brazil on May 25). We will report on this in the next

If your ILOC predictions lack some of the features that were in the EVANS predictions, you can request predictions that will at least have those features from your closest national or regional coordinator who has the PC version of the EVANS program (see below).

These predictions will be supplied free upon request for one station to IOTA members. Costs for extra stations, and for non-IOTA members, will be determined in a few weeks and announced in the next issue. If there are a large number of these requests, we may write the EVANS predictions to a magnetic tape and send it to ILOC for printing and distribution (as we did for many of the 1993 predictions), which would lower the cost for those from whom we received requests before finally writing the tape and sending it.

Overall, response to the 1993 prediction mailing was good, with we believe a higher percentage of returned forms than in other recent years. We have included in the updated datasets all of the verification forms that we received (and did not lose) before November 4. A second mailing, to those from whom we have no form, has not yet been made, but probably will be done later this month. Any responses to that mailing will be treated as updates, since they have been left out of the main run. However, for those requiring no changes, ILOC probably did generate the predictions (since they used the 1993 prediction dataset updates of a year ago for their main run), and ILOC can send them if we simply inform them of the address code.

PC Evans total occultation prediction software: The PC version of the EVANS program is working very well; the only problem is that a few relatively faint Kcatalog stars near the equator, involving less than 0.1% of the occultations, are treated as planets with unknown results in the predictions. Claudio Costa, from Rome, provided David with diskettes containing the executible program, Fortran source code, datasets, and some documentation at a small IOTA meeting held on Saturday, June 19, in Belgirate, Italy, just after the IAU Symposium, "Asteroids, Comets, Meteors 1993" (we thank Marco Cavagna for making arrangements for that meeting). Much of David's time since then has been devoted to documenting use of the program and its datasets, and managing the overall prediction process along the lines developed at USNO for their predictions generated with the mainframe EVANS program.

The first version of the "Evans package" was distributed at the ESOP-XII meeting in late August; see IOTA NEWS on p. 1. The documentation was finally completed in late October. The "Evans package" consists of much more than that program; it also includes the whole system of datasets, several small utility programs to manage them on a PC, and rather extensive documentation. Also included is a PC version of the LOCM program for generating local circumstances of asteroidal and planetary occultations, and software with documentation by Wolfgang Zimmermann for predicting the paths of occultations by

major and minor planets (a PC version of an early, but enhanced, version of David's MPOCC mainframe program). David Herald's OCCULT software, described on pages 306-307 of the last issue, is included an important note for those with OCCULT: install bat file with that program creates the 2-megabyte Watts file as a read-only file, so if you want to delete it later to make room for something else, you need to type the command "attrib watts.dat -r". We have deleted the "attrib" command at the end of install bat that makes watts dat read-only, and suggest that the other coordinators do the same, for future The Evans package consists of 27 distribution]. diskettes, most containing compact ZIPPED files. At least 60 megabytes of disk space are needed to use the software to generate predictions. The national and regional coordinators who now have, or who soon will have, the Evans package are listed below; addresses are in the IOTA-IOTA/ES Roster distributed with the last issue:

Graham Blow, New Zealand
Reinhold Buechner, Germany (IOTA/ES) *
Henk Bulder, the Netherlands
Claudio Costa, Italy *
David Dunham, Maryland *
Andrew Elliott, U.K.
Rick Frankenberger, Texas
Joaquim Garcia, Portugal *
Rocky Harper, Texas
Craig and Terri McManus, Kansas
Walter Morgan, California
Ton Schoenmaker, the Netherlands
Mitsuru Sôma, Japan
Matti Suhonen, Finland
Pierre Vingerhoets, Belgium

An asterisk (*) indicates that we have heard that the PC EVANS program has been installed and is ready to produce predictions. For operational use, subregions of the USNO super standard station regions have been defined and assigned to individual coordinators, so that the same station code is not used for different stations by different coordinators. Subregions will be assigned by David as he learns that individual coordinators get the PC EVANS program to work on their PC. If you have 60 megabytes to donate to the cause (at least on a temporary basis; the files can be backed up to diskettes, tape cartridge, Bernoulli disk, or other medium when the program is not in use) and are willing to provide predictions to IOTA members in your nation or region (IOTA can reimburse mailing costs for predictions sent to IOTA members), please contact David Dunham for a package. The only cost that we impose on receipients of the EVANS package is that they be willing to copy it and send it to one or two other coordinators whom we specify. In this way, Dunham and the other IOTA centers will not be spending all of their time copying and packaging diskettes. If you give a copy of the package to someone else, we want to know that, for distribution of new and updated datasets. With the widespread distribution of this software, IOTA can provide a prediction service that is locally more responsive than the previous centralized system has been, and we hope to get these predictions to all IOTA members who want them.

1994 Grazing Occultation Predictions: As you have seen in this issue and in other IOTA publications during the past year, Eberhard Riedel in Munich. Germany, has been preparing maps of grazing occultation paths for various purposes. Along with these maps, he also produces files of detailed graze limit predictions, an example of which is given at the bottom of this page. These predictions give essentially the same information as the current IOTA graze Riedel plans to provide packages to the IOTA graze computors that will include datasets of paths of grazes of stars to mag. 8.5 (also the USNO) limit) for each of the super standard station (SSS) regions [see the 1993 hemispheric graze supplement for their definition], and software that will read one of these datasets and the corresponding IOTA station file for the SSS region to generate detailed predictions of all grazes within each observer's favorable travel radius. For the rare (and usually spectacular) grazes of stars of mag. 4.5 and brighter (except in some cases near full moon when moonlight interference is too

strong), every observer in the region will get a prediction, regardless of their distance from the path. Riedel also plans to provide profiles for each graze, although this job could be done by the ACLPPP program. In any case, we plan to construct an interface with ACLPPP, so that we can maintain a consistent link with Mitsuru Sôma's OCCRED program that will be used also to analyze the observations. If Riedel's efforts to achieve the most accurate predictions (they already agree with the IOTA predictions to within about 200 m, usually smaller than the stellar and lunar limb errors involved) and the profiles delay too long the delivery of his packages to the graze computors. we may have difficulty getting predictions for January events to observers in time. For those few observers wanting predictions of 1994 "unfavorable" grazes not in Riedel's predictions (generally for stars fainter than mag. 8.5), David can use the OCC program following the same proceedures used for 1993 and earlier years. XZ Catalog: Both B1950 and J2000 versions of the USNO-IOTA prediction catalog of 32221 Zodiacal

stars, XZ80N (last updated in mid-1992), are now available from the Astronomical Data Center (ADC), National Space Science Data Center, Code 633, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA; telephone 1-301-286-8310 (Gail Schneider); fax 1-301-286-1771; Internet gail@ndadsa.gsfc.nasa.gov. Six other USNO catalogs that have been used for occultation predictions, including J, K, and L, are also now available from the ADC. Wolfgang Zimmermann is preparing a new version of the J2000 XZ catalog that will replace the positional data for most of the stars with more accurate data from the PPM catalog. I thank Wayne Warren for helping prepare the documentation for these catalogs.


# 39: JAN SOUTHERN DELTAT:	LIM	•	E	rar:		ONIS 138238, 1670,		6.4	1	57% SUNLIT, PHASE-ANGLE POS-AN.CUSE	E: 250.8
WEST N	ORT	H LATIT	JDE			MOON	MOON		SUN	POWI	CUSP
LONG.	DEG	. MIN.		7	JT	ALT.	AZI.	TANZ	ALT.	GRAZE	ANGLE
125.0000		36.30'			0.4s	48.3	195.8	0.89	-22.2	2 211.66	6.5 DA
124.0000		55.02'	13h	42m	36.2s	48.6	198.4	0.88	-20.7	7 211.67	6.5 DA
123.0000	36	13.65'	13h	45m	13.9s	48.8	201.1	0.88	-19.3	3 211.67	6.5 DA
122.0000		32.27'	13h	47m	53.0s	48.9	203.8	0.87	-17.8		6.5 DA
121.0000		50.99'	13h	50m	33.0s	48.9	206.5	0.87	-16.3	3 211.60	6.4 DA
120.0000	34	9.91'	13h	53m	13.7s	48.9	209.2	0.87	-14.8		6.4 DA
119.0000	33	29.15'	13h	55m	54.5s	48.8	211.9	0.88	-13.3	3 211.47	6.3 DA
118.0000	32	48.80'	13h	58m	35.0s	48.6	214.6	0.88	-11.7	7 211.37	6.2 DA
117.0000	32	8.96'	14h	1m	14.7s	48.3	217.2	0.89	-10.2	211.26	6.1 DA
116.0000	31	29.74'	14h	3m	53.1s	47.9	219.8	0.90	-8.6		5.9 DA
115.0000	30	51.22'	14h	6m	29.9s	47.4	222.3	0.92	-7.1		5.8 DA
114.0000	30	13.49	14h	9m	4.7s	46.9	224.7	0.93	-5.5		5.6 DA
113.0000	29	36.63'	14h	11m	37.0s	46.3	227.0	0.95	-4.0		5.5 DA
112.0000	29	0.71'	14h	14m	6.6s	45.7	229.2	0.98			5.3 DA

REPORTS OF ASTEROIDAL APPULSES AND OCCULTATIONS

Jim Stamm

If you do not have a regional coordinator who forwards your reports, they should be sent to me at: 11781 N. Joi Dr. Tucson, AZ 85737 USA. Names and addresses of regional coordinators are given in "From the Publisher" on Occultation Newsletter's front page. All times in this report are UTC.

Isao Sato has forwarded an observation that he recently came across: Miyoshi Ida reported an occultation of BD +25°1937 by (106) Dione on January 19, 1983 from Yokaichi, Japan. His report was made soon after the observation, but since there were no confirming observations made in Japan, the report was not accepted. Sato, recalling a 1984 article (Kristensen, L.K., Astron. Nachr. 305, p.207) about the event reported in Europe, used Ida's data and added the line to the diagram reproduced here.



I have summarized all of the reports that I have received for the first half of 1991 in the following two tables and section of notes. Table 1 lists the 1991 date, minor planet, occulted star, IDs of successful observers, and references to any notes. Table 2 lists the observer's ID, name, nearest town to location of observation, country (includes state or province for North America and Australia), and the total number of observations made in the period. The notes section details those events that included positive

observations, or other significant information that could not be reported in the tables. I am not including notes on those observations that may have been spurious unless there is some sort of confirmation, or the fact that something may have happened is relevant to another observation. Instead, I will place an asterisk (*) in the Notes column to indicate that I have received a report with more than a "no event....." in it.

Table 1. Asteroidal appulses and occultations; Jan.-June 1991,

1991	1	Min	or Planet	Cat	Star	Observers Notes
Jan	02	179	Klytaemnest.		21180	Hon
Jan	04	4	Vesta	AGK3	+10° 0317	1
Jan	09	185	Eun!ke	PPM	155732	MelBlkSmcAed
Jan		11	Parthenope	LickV	6202	Dss
Jar		184	De jope ja	PPM	97666	Så cAndMol
Jan	1.3	361	Myrrha	SAC	95912	2
Jar.			Oulu	₽₽M	93530	.3
Jan		11	Parthenope	LickV	1712	Dez
Jan	19	532	Her out I na	Licki	0954	Dez∂ta
Jan	1.9	216	Kleopatra	21 M	152432	4
Jan	19	1.7	Thet is	AGK3	491° 0500	Mol
Jan	21	230	Athamantis	SAO	156876	AvaCanDflGezGroKkn 5
						MosMtlBenTrlTfr
Jan	26	.94	Circe	LickV	30296	GroFl:ChsHozStwVie
						HellDwdDonBollBxlMcbGat
	33	27	Euterpe	PPM	126000	Sta
Feb	01	664	Judith	PPM	125047	LapDikBikNesPaj
						Andimo
Feb	0.2	5	Astraea	SAC	96089	StoAn iHawBlkPrl 6
						Bemisyrolik
Pern	04	201	Penelope	PPM	125971	DrizDebSci
Feb	1.5	187	Tamberta	SAC	158961	St a
Fec	22	360	Carlova	₽₽M	120671	AzaBqlBrtFauSrc
						lacMrxNelOspWtt
Mair	0!	627	Charis	₽₽M	126389	BdwGeyMrx8t h
Mair	0.4	535	Mont ague	LickV	8021	Mol
Mair	0.6	245	Vera	LickV	2005	LapBlk
1 6M	1.0	318	Magdalena	PPM	120542	Bff[sqlCvqFwlEoo *
						GenEffKhlMrxSzk
Mair	1.6	369	Aeria	ACK3	+26" (145	v _{TX}
Mair	1.7	₹34	Chicago	DAG.	1.90/723	Thin
Mair	1.7	(fC)	Echo	K°	20206	AnasmoHut.
Mar	20	121	Harmione	FAC	16/316	Thin
Mar	21	968	Petunla	PPM	177977	Smc
Mar	21	126	Joella	SAC	136589	Sinks
Mar	26	747	Winchester	PPM	128799	Hon
Mar	27	1117	Lamberta	SAO	183474	LapBlkKruNesAnd
Mar		56#	Princetonia	PPM	93204	BqsHlmHulDsmEq1Hff
						Krt MrxRthShtThoTvn
915	29	446	Tipporta	280	113344	Char
Matr	30	2	Pallas	EAC	118404	WC.
Mar	31	Œ,	Aegle	SAC	16875	Kbl
Aur		217	Eludora	PPM	179112	StigBukAndButDEk 1
Apr		624	Hektor	SAO	181911	Cos Lea Worlden
Az:r		683	lanzia	(14)	186127	And
Apr		17	Thetis	Licky	23742	Wir
Apr		532	Hermulina	FAC	250.938	vázt.
Apr		184	De jope ja	AC	5827	Henlicz
Apr		532	Hermil (na	FAC	255192	BblDssdur.
Apr		4	Vest.a	AGK3	+20° 0416	St a
Ap:	11	121	Hermione	FAC	199379	NesBlkHLtAndGer
	13	19	Fortuna	AC	4776	GkoMea5c1Zwk
	15	177	Irma	SAO	184383	BilborOveDebdooMae
				·		Prosmi GadTim
Apr	15	270	Anabita	LickV	972	Sta
Apr	19	121	Remione	FIAC	211278	DebSmjOve(rr
Apr	19	103	Hera	LickV	12536	DikatqAndBik
Apr		175	Thyra	Yale	9594	BakGendildubica
Apr			Celuta	SAO	208561	Bik
May		15	Eunomia	EAC	368210	
лау Мау		207	Hedda	SAO	7986B	MolMiaHutKeaHemSmcAnd
. na y		701	resultad	SMC	1 2000	MiaAnd

Table 1 (Cont.). Asteroidal appulses/occultations: Jan.-June 1991.

1991 Minor Planet		Cat	Star	Observers	Notes	
May 08	462	Eriphyla	SAO	158878	Diz	
May 08	1345	Potomac	SAO	164518	St.gB1kDa1	
May 10	181	Eucharis	SAO	144646	StgSmcAndLoa	
May 11	102	Miriam	AC.	23399	Sta	
May 11	1655	Comas Sola	PPM	128119	DizSta	
May 24	1404	Ajax	SAO	181037	NesSmc	
May 27	674	Rachele	SAO	184425	EnkF1eOveDebThn	*
					BrrFnzGez	
Jun 08	585	Bilkis	PPM	174544	St.gSmc	
Jun 09	146	Lucina	FAC	447154	And	
Jun 11	694	Ekard	SAO	158489	Fga	
Jun 11	433	Eros	5 A O	226005	SmcHutAndSugfakIk	a
Jun 13	423	Diotima	SAO	185272	DikAndLoa	8
Jun 15	356	Liguria	SAO	210543	LunDebSmiThn	9
Jun 16	130	Elektra	PPM	168528	Sta	
Jun 19	776	Berbericia	SAO	159636	Smi Thn	
Jun 24	103	Hera	₽₽M	159359	CopOveSmi	
Jun 24	514	Armida	PPM	156789	CopOveSmi	
Jun 24	978	Aidamina	PPM	163619	Hon	
Jun 25	53	Kalypso	SAC	164279	Grc	
Jun 26	41	Daphne	ACK3	+06° 0115	OveSm1VnbThn	
Jun 29	897	Lys Ist rata	P₽M	171718	SmcAnd	

Table 2. Observers and locations of events Jan.-June 1991.

<u>I ab</u>	ie Z. Observers an	id locations of	events janjune 15	<u>/21</u>
ID	Observer	Town	Country	No
Alk	Aikman, Chris	Victoria	Brit. Columbia - CAN	1
Axa	Akazawa, Hidehiko	Funaho	Japan	1
Aki	Akiyoshi, T.	Miyazaki	Japan	1
Alw	Allis, Tanner	Midland	Michigan - USA	1
And	Anderson, Peter	The Gap	Queensland - AUS	18
Aza	Azema, JM.	Lama lou	France	2
Blg	Ballegoy, E.V.	Druten	Netherlands	1
Ban	Ban, Yoshihiko	Mastumoto	Japan	2
Bff	Baruffetti, Pietro	Massa	Italy	,
Bay	Bayus, Chris	F11nt	Michigan - USA	:
Bem	Bembrick, Colin	Bathurst	N.S.W AUS	2
Brilw	Benedyktowicz, L.	Kracow	Poland	
Bgl	Bertoglio, A	Torino	Italy	
Brt	Bertoli, Oreste	Alpignano	Įtaly	7
B≭v	Bettonvil, F.	Heesch	Netherlands]
Bzl	Binzel, Richard	Lexington	Massachusetts - USA	1
Bis	Blshop, Roy	Avonport	Nova Scotia - CAN	1
Blk	Blanksby, Jim	Wandin	Victoria - AUS	12
Blm	Blommers, L.	Leiden	Netherlands	
Bst	Holster, Hob	Alexandria	Virginia - USA	
Biam	Boltman, A. & M.		South Africa	-
Bol	Boltwood, Paul	Ottawa	Ontario - CAN	
Brr	Borras, Vincente	Benicarlo	Spalt:	
Ek 25	Boschat, Michael	Halifax	Nova Scotia - CAN	
9gs	Bourgeols, Jean	Ciney	Belgtum	
Bb)	Briet Blanes, T.	Castellon	Spain	
Bri	Brisbin, Jack	Dryden	Mlchlgan - USA	
Bru	Bruce, Robert	Drouin	Victoria - AUS	
Bul	Bulder, Henk	2oetermeer	Netherlands	
Bun	Burns, Tom	Stratford	Ohio - USA	
Вут	Byron, Jeff	Sydney	N.S.W AUS	
Can	Candela, Bernard	Sollies-Pont	France	
Car	Carlson, Jim	Falmouth	Massachusetts - USA	
Cas	Casas, Ricard	la Orotava	Spain	
Cat	Castle, Paul	Colona	Illinois - USA	
Cvq	Cavagna, Marco	Sormano	Italy	
Chs	Chase, Rick	S. Waterford	Maine - USA	
Cir	Cirker, Willkie	Lucas	Ohio - USA	
Copp	Cooper, Tim	Benoni	South Africa	
Cos		Coal Valley	lilinois - USA	
Cun	Cunningham, M.	Whittier	California - USA	
Dal		Launcet on	Tasmania	
Df1	Daiffallah, Khalil	Alger	Algeria	
Dae	· ·	etermaritzburg	South Africa	
Deab	•	Ladysmith	South Africa	
Dnz	Denzau, Helmut	Essen	Germany	

New Zealand

Table 2 (Co	ont.). Obser	rvers/location:	s of events:	Jan-June 91

ID	Observer	Town	Country	No
Did	Didick, Richard	Norton	Massachusetts - USA	1
Diz	Dietz, Richard	Greeley	Colorado - USA	2
Dod	Dodson, Steve	Sudbury	Ontario - CAN	1
Dom	Dombrowski, Phil	Glastonbury	Connecticut - USA	1
Dow	Dow, Donaldson	Ramrod Key	Florida - USA	1 1
Dwd Dwd	Dunham, David W.	Greenbelt	Maryland - USA New Jersey - USA	1
Dw2	Dunham, David W. Dunham, David W.	Maryland Detroit Air.	Michigan - USA	1
Dw3 Dss	Dusser, Raymond	Kalaa Sghira	Tunisia	5
Equ	Egawa, Humiharu	Kumatori	Japan	:
Eir	Eiraku, M.	Uchinoura	Japan	1
E11	Elliott, A.J.	Reading	United Kingdom]
Enm	Emmons, Richard	North Canton	Ohio - USA	1
Enk	Enke, S.	Windhoek	South Africa	:
Ew1	Ewald, D.	Blesent hal	Germany	1
Fau	Faure, G.	Varces	France	1
Fnz	Fernandez B., D.	Barcelona	Spain	1
Fle	Field, R.	Durban	South Africa	1
Fit	Fletcher, Murray	Victoria	Brit. Columbia - CAN	
Foc	Fodor, L.	Kunadads	Hungary	1
F.dJ	Foglia, Sergio	Lacona	Italy	1
Ex)	Fox, Jim	Afton	Minnesota - USA	1
Frs	Frasca, Michael	Peoria	Illinois - USA	
Fry	Frey, Gary	W Bloomfield	Mlchigan - USA	1
Fuj	Fujiwara, Y.	Osaka	Japan	1
Fuk	Fukushima, S.	Machida Lisboa	Japan Doctors	1
Gre	Garcia, Joaquim		Portugal South Africa	
Gad Gat	Garde, T. Garrett, Lawrence	Shurugwe Burlington	Vermont - USA	
Сел Сел	Genovese, Marco	Torino	Italy	
Gern	George, Martin	Launceton	Tasmania	- :
Ger	Gerry, Karen	Jones Beach	New York - USA	3
Gib	Giles, Barry	Hobart	Tasman!a	7
Gli	Gill, David	Massilon	Ohio - USA	7
Ggr	Gomez de Quiroga	Madrid	Spain	3
Gez	Gomez, F.J.	Mollet	Spain	
Gov	Gondalves, Rui	Lisboa	Portugal	
Gko	Gorko, M	Tzxtz	Poland	
Grh	Graham, Frances	E. Pittsburg	Pennsylvania - USA	
Gro	Grainger Obs.	Exeter	New Hampshire - USA	
Grs	Grieser, Dan	Columbus	Obto - USA	•
Gur	Guerber, Jeff	Greenbeit	Maryland = USA	
Cry	Suyer, E.H.	Daun	Germany	
Bas	Hasegawa, Takashi	Tochiqi	Japan	
Hsh	Hashimoto, Akie	Chichiba	Јараг:	
Hat.	Hattori, Y.	Miyazaki	Japan Taran	
Haa TVti	Hayashi, H.	Zushi	Japan United Kingdom	
	Haymes, Tim V. Hays, Robert	Reading Worth	Illinois - USA	
Hay Haw	Hayward, Steve	Madang	Papua New Suinea	
llel	Heil, Christopher	Andover	Massachusetts - USA	
Henri		Brooklyn	Wisconsin - USA	
Hey	Henry, Michael	Cant on	Ohio - USA	
Hed	Heydon, Alex	Toronto	Ontario - 'AN	
Hí!	Hill, Kym	Hobart	Tasmania	
Hir	Hirose, T.	Ota	Japan	
lil s	Hirose, Y.	Chlgasaki	capan	
Hff	Hoffman, Martin	Weidenbach	Germany	
Hoz	Holtz, John	Russellton	Pennsylvania = USA	
Barn		Fuchu	Japan	
Hon	Honkus, Edward	Carnegie	Pennsylvania - USA	
Hor	Horiuchi, J.	Ohlto	Japan	
Hos	iloso!, S.	Soka	Japan	
Hut	Hutcheon, Steve	Sheldon	Queensland - AUS	
lac	Lacovone, N.	Torino	Italy	
]ka	Tkari, Yasukazu	Taga Taga	Japan Japan	
Ima	Imal, Y.	Toshima	Japan South Africa	
Joo		Reitz	South Africa Minnesota - USA	
Kac		Roseville	Indiana - USA	
Kai Kak	· ·	Columbus Shizuoka	Japan	
Kan		Sakura	Japan Japan	
Kar		Nerima	Japan	
Kesa		Bundaberg	Queensiand - AUS	
Ket	Keith, Lee	Milwaukee	Wisconsin - USA	

Tab	ole 2 (Cont.) Obse	rvers/location	s of events: Jan-June	• 91
ID	Observer	Town	Country	N
Kel Kit	17	Halifax	Nova Scotia - CAN	
		Musahino	Japan	
КijУ		Zama	Japan	
Kom		Wald	Switzerland	
Kkn		Sakado	Japan	
Koy	,	Salonta	Romania	
Krt	- •	Hachioji	Japan	
Kru	, ,	Siegen	Germany	
Kuj		,	Victoria - AUS	
Kum		West Alits	Wisconsin - USA	
Lac		Kumamoto	Japan	
Iap		Halifax	Nova Scotia - CAN	
Lgn		The Basin	Victoria - AUS	
Lea	,	Villeneuve	France	
Jlx		East Rand	South Africa	
lez	•	Meudon	France	
Ind		Wilton	Connecticut - USA	
Ioa		Esbjerg Christchurch	Denmark	
Low	•	Calgary		
Lun	•	Johannesburg	Alberta - CAN	
Lur	• •	West Chester		
Modo		Bedford	,	
Mae	,	Klyotake	Massachusetts - USA	•
Mal	Moley, Paul	Metamora	Japan Michigan uga	
Man		Tempe	Michigan - USA Arizona - USA	
Mas		Brooklyn	Wisconsin - USA	
Mrc		Bloomington	Indiana - USA	
Mit	Marlot, C.	Guines	France	
Mt.1	Marti, Josep	Mataro	Spain	
Mrx		Stuttgart	Germany	1
Mico		Olta	Japan	
Mat	Matsuda, Hideki	Tenri	Japan	
Mem		Falmouth	Massachusetts - USA	
M:w	McWatters, Richard		Ontario - CAN	,
Mia	Miller, Anderson	Jarrahdale	W. Australia - AUS	
Min	Minobe, J.	Takasaki	Japan	2
Miz	Miyazaki U. Ast Ch	ib Kivotake	Japan Japan	١
Mod	Modic, Robert	Richmond Hts		1
Mo:	Moller, Harry	Kingsley	W. Australia - AUS	2
Моу	Mortya, Y.	Yokohama	Japan	,
Mos	Mostefaoul, Toufik	Alger	Algeria	2
Mur_	Muto, N.	Sakado	Japan	1
Ne1	Neel, Regis	Venissieux	France	1
Nes	Nelson, Peter	Ellinbank	Victoria - AUS	4
Ohp	Obs Haute-Provence		France	1
Jun	Obs. Jungfrauloch	Jungfraujoch		1
Metu		Meudon	France	- 4
Ona	Odaglri, S.	Aomori	Japan	1
Ohk	Okura, Nobuo	Okavama	Japan	1
Clv	Oliver	Bronson	Florida - USA	1
Osb	Osborn, Wayne	Mt. Pleasant	Michigan - USA)
Œh	Oshima, O.	Kurashiki	Japan	1
Ove	Overbeek, Danie	Edenvale	South Africa	8
Oya	Оуа, к.	Chiqasaki	Japan	1
Pa.	Paller, Leroy	Central	Arizona - USA	1
Paj	Park, J.	Melbourne	Victoria - AUS	2
Uck	Peorink, U	Heesch	Netherlands	i
Per	Persha, G.	Lowell	Michigan - USA	1
Pe 1	Peterson, John	Falmouth	Massachusetts - USA	1
Pek	Petrasko, Michael	Chat ham	Massachusetts - USA	1
Ph1	Phillips, Jim	Grand Rapids	Minnesota - USA	1
Plu	Pluth, Bernard Jr.	Dublin	Ohio - USA	1
Poa	Poani, Ernie	amphell's is.	Illinois - USA	1
Pr1	Priestley, John	Pukerua Bay	New Zealand	1
Pro		termaritzburg		1
Pry	Pryal, Jim	Seattle	Washington - USA	1
Ros	Ross, Hob	Florence	Kentucky - USA	1
Rt.h	Rothe, Wolfgang	Berlin	Germany	3
Sao	Sato, I.	Meguro	Japan	1
Sao	Sato, M.	Fuchu	Japan	1
Sci	Scarfi, G.	La Spezia	Italy	1
Sh:	Schaaf, Elizabeth		Iowa - USA	3
Scto	Schnabel, Carles	Barcelona	Spain	3
Sht.	Scholten, Alex	Eerbeek	Netherlands	1
				•

Table 2 (Cont.). Observers/locations of events: Jan-June 19						
ID	O DUCI TO	Town	Country	N		
Scm		St.rat.ford	Ohio - USA			
Shb		Matsubarako	Japan			
Shm	Shima, K.	Fujimi	Japan			
Sml	Small, Craig	Jones Beach	New York - USA			
Smi	Smit, J_	Pretoria	South Africa			
Smc	Smith, Charlie	Woodridge	Queensland - AUS	14		
Smg	Smlth, Greg	Elyrla	Ohio - USA			
Som	Soma, M.	Mitaka	Japan			
Stg	St. George, Lou	Auckland	New Zealand			
Sta	Stamm, Jim	Tucson	Arizona - USA			
St.h	Storch, Samuel	Jones Beach	New York - USA			
St₩	Stowers, Clifford	Fairfield	Maine - USA			
Sun	Suenaga, Y.	Fulinomiya	Japan			
Sga	Sugai, M.	Aikawa	Japan			
Sug	Sugie, J.	Taga	Japan			
Suz	Suzuki, A.	Yamato	Japan			
Suk	Suzuki, Y.	Kodaira	Japan			
Szk	Szarka, Levente	Kecskemet	Hungary			
Tak	Takahashi, S.	Taga	Japan			
Tan	Tanaka, K. His	gashimatsuyama	Japan			
Tr	Terrier, Pierre	Chamonix	France			
Thn	Thompson, R.		South Africa			
Ths	Thompson, Ray	Toront o	Ontario - CAN			
T'ho	Thooris, Bertrand	Wervik	Belgium			
Thh	Thrush, Jeff	Taylor	Michigan - USA			
Tky	Tokyo U. Ast Club	Meguro	Japan			
Torn	Tomoda, T.	Kobayashi	Japan			
Tor	Torihara, Y.	Miyazak:	Japan			
Trl	Torrell, Sebastia	Barcelona	Spain			
Uexi		Habikino	Japan			
Vnb	VanBlommestein, P.	Simon's Town	South Africa			
Fvl	Van Loo, Francois	Genk	Belgium			
Van	Vance, Norbert	Allen Park	Michigan - USA			
V!f	VanEffen, Dick	Midland	Michigan - USA	-		
Vit	Victor, Robert	E. Lansing	Michigan - USA	,		
Vie	Viens, JF.	Quebec City	Quebec - CAN	-		
Waa	Wagenaar, R	Wateringen	Netherlands	2		
Wku	Waku, S.	Kat sushika	Japan			
Wrr	Warren, Wayne	Maryland	New Jersey - USA	_		
White	Watanabe, O.	Sapporo	Japan	-		
Yat.	Watanabe, Shinobu	Saita	Japan			
Wei	Weier, David	Brook!yn	•			
We !	Welch, Douglas	Hamilton	Wisconsin - USA Ontario - CAN			
Wld	Wild, Walter	Chicago				
WII	Wilds, Richard	•	Illinois - USA			
Witt	Witt, U.	Topeka	Kansas - USA	2		
Yam	Yamada, M.	Berlin	Germany	2		
Yur	Yurchesyn, Joe	Yokohama	Japan	:		
Zwk	Zawilski, Marek	Halifax	Nova Scotia - CAN	1		
OMA	nowsiaki, marek	Lodz	Poland	1		

NOTES:

- Jan ()4 (4) Vesta. See [O.N. 5(12), p.315]. Also see Jan. '92 [Sky & Telescope 83(1), p.72]. Positive observations came from HedThsMcwDw3PovFryBri ModHozOsbMal PerPluSmgRosGllAlwVffEmmGrsDod VanThh. Negatives from DssMosMtiTrlMrcHonHay PhlBosKelLae YurDidFrsDowKacBun ScmWilWasWei HnnSamKujVitMacDunOlv KaiKeiBayBzlVicBisCatPoa ShfCosWld.
- 2 Jan 13 (381) Myrrha. See [O.N. 5(12), p.315]. Isao Sato, Mitsuru Soma, and Toshio Hirose have authored The Occultation of Gamma Geminorum By the Asteroid (381) Myrrha in the Apr. '93 [Astronomical Journal 105(4), pp.1553-61]. Sato (in a communication to IOTA) reports three additional video events not included in the AJ. paper. One was an 8.35

second occultation beginning at approximately 11:59:54 from Kiyosato, Japan (Longitude: 138"24'28"E. Latitude: 35°55'59"N. Height: 1500 meters), by Akira Ogino. The companion of Gamma Geminorum can be seen throughout the event. Sato explains that this fact is inconsistent with the visual observation of its emersion after the disappearance of the primary by Shu Kaneko, who was nearby on the occultation track. One interpretation of this inconsistency is that the brightness of the primary delayed Kaneko's recognition of the companion. If so, the position of the companion should be modified. Another interpretation is the existence of a third body, fainter than the companion.

This was the brightest and best observed event ever seen in Japan of an asteroidal occultation. 32 chords from 54 observers were obtained. Myrrha was shown to be an ellipse of about 147 by 127 km. Gamma Geminorum was also shown to have an upper limit in diameter of .0026 arcsec. Its 7.5 magnitude companion is separated by .06 arcsec. Positive observations came from HasTanBanKomMutShbHshKanHosWkuKarShmImaKit TkySaoSom SsoHomKoyHirMoy FukSuzSgaYamKiy OyaHisHaaSunHorOgi. Negatives from WnbOdaSuk KakIdaWatFui UedOhkAkaOshMto KumHatTorAkiTom MaeMizMinEir. There were 4 positive observations in China (along with about 3000 misses!) - not yet reported. The Royal Astronomical Society of New Zealand also reports negative observations from KruBruPajBlkLapAndSmcHut.

- 3 Jan 13 (1512) Oulu. Observers were BgsBlgBlmBtv BulCasDnz DssEllFvlGev GeyGqrGrcJlxLgn LhdMltRth TrlTvhUpkWaa.
- 4 Jan 19 (216) Kleopatra. See [O.N. 5(12), p.315]. Also see Jan. '92 [Sky & Telescope 83(1), p. 72]. Positive observations came from Dw2WrrGurAikMod SthPryLurHey. Negatives from McbGrsGroCirBunChs CunFttCarPej MemPekGatBstGer SmlVieManPalMis LowLezFxj. A blink of less than a second duration was reported by Wil.
- 5 Jan 21 (230) Athamantis. See [O.N. 5(12), p.315], and a separate article in this issue. Positive events were reported by AzaDfiGezMosMtiScbTrl. Negatives by CanGrcKknTrr
- 6 Feb 02 (5) Astraea. See [O.N. 5(12), p.315].
- 7 Apr 02 (217) Eudora. See [O.N. 5(12), p.315].
- 8 Jun 13 (423) Diotima. Sec [O.N. 5(12), p.315].
- 9 Jun 15 (356) Liguria. See [O.N. 5(12), p.315].

Several observers have indicated that the long time span between observation and publication tends to discourage making the observation and reporting it. I agree, and from now on I will publish summaries in a more timely fashion, from the reports that I have on hand. Reports received after my summary is written will be archived but not published.


Of course this does not apply to positive observations that require the collection and reduction of data.

Edwin Goffin reports some changes to his 1994 asteroidal occultation predictions based on improved or erroneous elements. Some changes move the path a hundred or so miles, but errors generally shift the path that much anyway, so only the significant updates follow, including newer world maps from Goffin.

Discard predictions for (338) Budrosa, (399) Persephone, (611) Valeria, and (476) Hedwig. Add the events for Apr 17, Jun 17, and Oct 19. Note time and/or path changes for the events on Feb 04, Jun 28, Oct 30, and Nov 26.

[Ed. The additional events will not be visible from populous parts of North American, so copies of them are being sent to the appropriate regional coordinators for distribution rathern than being published here. For the changed events, we include only Goffin's titles and world maps, since the other information on his chars is not changed.]


[Unfortunately, there is not enough time to include the article on the 1991 January Athamantis event mentioned in Jim's article. We plan to include it in the next issue, and some information about it will be given in the 1994 February issue of Sky and Telescope.]




326 Tamara - PPM 52402 1994 feb. 4 9h52.7m U.T. \$40 43724




2393 Suzuki — PPM 233040 1994 jun 28 9h 9.9m U.T.





499 Venusia — PPM 270020 1994 oct 30 18h13,0m U.I.




514 Armida — PPM 117739 1994 nov 26 - 3ht8.9m U.I.









SOLAR SYSTEM OCCULTATIONS IN 1993

David W. Dunham

This is a continuation of the article on pages 315-320 of the last issue, which in turn was started on p. 257 of ON 5 (10). Regional maps showing asteroidal and planetary occultation paths for November and December, similar to those in ON 5 (10) for February through May, are included in this issue. Unfortunately, there was not time to prepare finder charts for events that were not included in Edwin Goffin's predictions (with finder charts) that were distributed over a year ago.

There are many good events during the last two months of 1993, especially in North America and Europe. The usual world maps produced by Mitsuru Sôma (using my data for the occultation) are included for many of them. I have drawn Edwin Goffin's predicted paths on these if they differ from my prediction by more than 0".5. I have also put "PPMh" after "GOFFIN" on the label for these paths if Goffin has used positional data for the star from the especially accurate high-precision subset of the PPM catalog. If Goffin's time differs from my calculated time by more than 4 minutes, I have written the difference, Goffin's time minus my time, in minutes on the upper left or upper right side of the world figure. For example, "+9"" means that Goffin predicts that the time of closest approach for any location will be 9 minutes later than the time of closest approach calculated with my data and given in the local circumstance predictions distributed by Joseph Carroll and by IOTA/ES.

Notes about Individual Events:

July 3: I distributed a catalog of relatively accurate positions of stars to about 11th magnitude from the Lick Voyager-Uranus (LU) catalog near the path of 24 Themis to potential astrometrists. Astrometry obtained at Climenhaga Observatory showed a small south shift of this path, so that it probably extended from Washington state and southern British Columbia across Montana, the Dakotas, and part of Minnesota, but known observers in this path were clouded out.

Aug. 18, Fortuna: SAO 163155 is in the Lick Voyag er-Neptune catalog, so again I was able to distribute a subcatalog of fainter stars along the track to astrometrists. Alan Gilmore obtained plates on two nights at Mt. John Observatory in New Zealand. Analysis of measurements of these plates showed that the path shifted a small distance southward, still over Cuba and the Bahamas. J. Allyn Smith, Melbourne, FL, confirmed that the asteroid passed south of the star at the updated time, as expected from the astrometry.

Oct. 3, Pluto: Lick Observatory CCD astrometry of Pluto and the target star late in September showed that the path of this occultation missed the Earth's surface to the north. Based on this information, international expeditions to observe the occultation were cancelled. The astrometry showed that, within the uncertainties, an occultation by Charon was possible in either New Zealand or (less likely) eastern Australia. Graham Blow reports that the appulse was monitored photoelectrically at Mt. John Obs. through some cirrus at low altitude, and no occultation was detected.

Oct. 9, Euterpe: Astrometry obtained at Lowell Observatory on Oct. 5 showed that the nominal path was quite accurate, although there was a sizeable correction to the time, 6 min. earlier than my prediction. Unfortunately, all observers in the path from Kansas eastward were clouded or fogged out. Some observers in Arizona also had problems with clouds, but several there, including Jim Stamm, timed the occultation. Observations were also made in New Mexico and by Paul Maley with an image-intensified video system at Dumas, in the Texas panhandle. Two other intensified video recordings were made, at Belen, Albuquerque and at Central Arizona College Observatory in Casa Grande, where a 2-second occultation near the northern limit occurred. Albert Vreeland, at his home in southern Tucson, AZ, had only a quick dimming during the correct second, marking the southern limit. I believe that three photoelectric records were obtained. Since the maximum durations were less than 11 seconds, Euterpe's outline may have been a little smaller than predicted. The actual occultation path is plotted on the Western Hemisphere regional map. More about this event will be given in a future issue.

Noy. 7, Chiron: This event was first publicized only two months ago and has not appeared in previous ON's. It was identified by S. Bus, L. Wasserman, and J. Elliot in a special search of occultations by this distant comet-like object during 1993-1996; a paper about them, including small finder charts, has been submitted to Astron. J. Around 13:20 UT of Nov. 7, Chiron was predicted to occult 14.3-mag. GSC 249-00971 along a path crossing the southwestern USA. It was the only event they found that had a reasonable chance of being visible from the contiguous USA. A special notice including part of the preprint giving information about this event was mailed to IOTA members in the possible region whom we thought might have access to large telescopes.

Nov. 13, Apollonia: The star is ZC 3286

Nov. 20: 288 Glauke has a very long-period light curve, thought possibly due to precession caused by a satellite of the asteroid. But radar observations do not

show two objects, only one very slowly rotating one. Nov. 23, Fortuna: A major effort for this event is planned, similar to the one for Euterpe in early October. Observers in the southern and midwestern USA, especially in the region between (and slightly north and south of) my and Edwin Goffin's paths, are encouraged to provide me with longitudes, latitudes, and heights above sealevel, preferably accurate to a km or less, of all known sites where observations might be obtained. The computerized sort of stations relative to the updated path used for the Oct. Euterpe event can be used again for this event, and the results distributed by E-mail, for targeting of mobile observers to gaps in the fixed-site coverage of the event.


Nov. 23, Crocus: Like Glauke, 1220 Crocus has a very long-period light curve. A satellite may be responsible; there are no radar observations of Crocus. Nov. 29: This event occurs during the total lunar eclipse discussed earlier in this issue. The star is ZC 2025. Wolfgang Beisker and Hans Bode plan to obtain a CCD record of the occultation from an observatory in the Canary Islands.


Dec. 11, Victoria: The star is ZC 466.


Dec. 20, P/Schwassmann-Wachmann-1: This giant comet beyond Jupiter's orbit may occult 13.1-mag. GSC 19150293 somewhere in eastern North America and Europe around 1:50 UT, according to new calculations by Edwin Goffin; the event, and two other occultations that he found, are not in any of IOTA's previously-distributed material. A copy of Goffin's chart for either of these events is available upon request to either the McManuses in Topeka (phone 1-913-232-3693, E-mail 570-0611@mcimail.com, address given at top of p. 1) or to Hans Bode (see back page). The 2nd event is of 13.3-mag. GSC 19150001 around 13:50 UT Dec. 23, possible area Siberia, Japan, and western North America; and the 3rd event is of 13.4-mag, GSC 19150481 around 7:00 UT Dec. 25, possible area North America, northwestern Africa, and Europe from Iberia to Scandinavia.


<u>Dec. 21, Metis:</u> Special efforts are planned to update the path prediction for this good event for Australia or New Zealand, possibly using observations of past occultations by Metis.


Dec. 31, Vibilia: The star is possibly a close double star, based on a "gradual" lunar occultation observed in South Africa on 1928 April 24.























The International Occultation Timing Association was established to encourage and facilitate the observation of occultations and eclipses. It provides predictions for grazing occultations of stars by the Moon and predictions for occultations of stars by asteroids and planets, information on observing equipment and techniques, and reports to the members of observations made. IOTA is a tax-exempt organization under section 509(a)(2) of the (USA) Internal Revenue Code, and is incorporated in the state of Iexas.

The ON is the IOTA newsletter and is published approximately four times a year. It is also available separately to non-members.

The officers of IOTA are:

President David W. Dunham Executive Vice President Paul Maley **Executive Secretary** Rocky Harper Craig and Terri McManus Secretary-Treasurer for Grazing Occultation Services Joe Senne VP for Planetary Occin Services
VP for Lunar Occultation Services Joseph Carroll Walter Morgan ON Editor Joan Bixby Dunham Hans-Joachim Bode 10TA/European Section President IOTA/ES Secretary IOTA/ES Treasurer Eberhard Bredner Alfons Gabel IOTA/ES Research & Development Wolfgang Beisker Eberhard Riedel IOTA/ES Public Relations

Addresses, membership and subscription rates, and information on where to write for predictions are found on the front page.

The Dunhams maintain the occultation information line at 301-474-4945. Messages may also be left at that number. When updates become available for asteroidal occultations in the central U.S.A., the information can also be obtained from either 708-259-2376 (Chicago) or 713-488-6871 (Houston).


Observers from Europe and the British isles should join IOTA/ES, sending DN 40,-- to the account IOTA/ES; Bartold-Knaust Strasse 8; 3000 Hannover 91; Postgiro Hannover 555 829 - 303; bank-code-number (Bankleitzahl) 250 100 30. Full membership in IOTA/ES includes the supplement for European observers (total and grazing occultations) and minor planet occultation data, including last-minute predictions, when available.

The addresses for IOTA/ES are:


Eberhard Bredner Ginsterweg 14 Bartold-Knaust-Str. 8 D-59229 Ahlen 4 (Dolberg) Germany Germany

Telephones (49- for Germany, -2388-3658 -511-424696


Fax -2381-36770 Hans-Joachim Bode Bartold-Knaust-Str. 8 D-30459 Hannover Germany or 0 within Germany) -511-424696




SAO 128628 by Metis 93 Dec 21




SAO115514 by Klotho 93 Dec 22 SAO 79240 by Pompeja Dec 26



SAO 109579 by Parthenope Dec 31



Anonymous by Crocus Dec 31



SAO 78468 by Vibilia 93 Dec 31